

Datasheet for ABIN7467961

anti-APEX1 antibody



Go to Product page

_			
()	11/0	r\ /	iew
	' V C	IV	I C. V V

Quantity:	100 μL
Target:	APEX1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This APEX1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	Full length human APE1 purified from E. coli
Clone:	2104
Isotype:	lgG1
Cross-Reactivity:	Human, Mouse
Purification:	Protein G purified

Target Details

Target:	APEX1
Alternative Name:	apurinic/apyrimidinic endodeoxyribonuclease 1 (APEX1 Products)
Background:	Synonyms: apurinic/apyrimidinic endodeoxyribonuclease 1 , APE , APE1 , APEN , APEX , APX , HAP1 , REF1

Background: Apurinic/apyrimidinic (AP) sites occur frequently in DNA molecules by	
spontaneous hydrolysis, by DNA damaging agents or by DNA glycosylases that remove specific	
abnormal bases. AP sites are pre-mutagenic lesions that can prevent normal DNA replication	
so the cell contains systems to identify and repair such sites. Class II AP endonucleases cleave	
the phosphodiester backbone 5' to the AP site. This gene encodes the major AP endonuclease	
in human cells. Splice variants have been found for this gene, all encode the same protein.	
[provided by RefSeq, Jul 2008]	

Molecular Weight:	36 kDa
Gene ID:	328
UniProt:	P27695
Pathways:	DNA Damage Repair, Chromatin Binding, Cell RedoxHomeostasis, Smooth Muscle Cell
	Migration, Positive Regulation of Response to DNA Damage Stimulus

Application Details

Application Notes:	WB: 1:1000. IHC-P: 1:5000. ELISA: 1:1000-1:10000. Optimal dilutions/concentrations should be
	determined by the researcher. Not tested in other applications.
Comment: Positive Control: HepG2, HeLa, Raji whole cell lysates	
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.2, No Preservative
Preservative:	Without preservative
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.