

Datasheet for ABIN7468115 **anti-CD3G antibody (C-Term)**



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	CD3G
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CD3G antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	Carrier-protein conjugated synthetic peptide encompassing a sequence within the C-terminus region of human CD3 gamma. The exact sequence is proprietary.
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by antigen-affinity chromatography.

Target Details

Target:	CD3G
Alternative Name:	CD3g molecule (CD3G Products)

Target Details

Background:	Synonyms: CD3g molecule , CD3-GAMMA , IMD17 , T3G Background: The protein encoded by this gene is the CD3-gamma polypeptide, which together with CD3-epsilon, -delta and -zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T-cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. Defects in this gene are associated with T cell immunodeficiency. [provided by RefSeq]
Molecular Weight:	20 kDa
Gene ID:	917
UniProt:	P09693
Pathways:	TCR Signaling , CXCR4-mediated Signaling Events

Application Details

Application Notes:	WB: 1:500-1:3000. IHC-P: 1:100-1:1000. FACS: 1:50-1:200. Optimal dilutions/concentrations should be determined by the researcher. Not tested in other applications.
Comment:	Positive Control: Molt-4
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.1M Tris-Glycine (pH 7), 10 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.