# antibodies -online.com







## anti-LIN7A antibody



| ( ) | ve     | K\ / |    | A . |
|-----|--------|------|----|-----|
|     | $\cup$ | 1 V/ | Щ. | V۷  |
|     |        |      |    |     |

| Quantity:    | 100 μL  |
|--------------|---|
| Target:      | LIN7A   |
| Reactivity:  | Human   |
| Host:        | Rabbit  |
| Clonality:   | Polyclonal  |
| Conjugate:   | This LIN7A antibody is un-conjugated  |
| Application: | Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Whole Mount) (IHC (wm)) |

#### **Product Details**

| Immunogen:        | Recombinant protein encompassing a sequence within the center region of human LIN7A. The exact sequence is proprietary. |
|-------------------|---|
| Isotype:          | IgG   |
| Cross-Reactivity: | Human, Mouse, Zebrafish (Danio rerio)   |
| Purification:     | Purified by antigen-affinity chromatography.  |

## Target Details

| Target:           | LIN7A  |
|-------------------|--|
| Alternative Name: | lin-7 homolog A, crumbs cell polarity complex component (LIN7A Products)                     |
| Background:       | Synonyms: lin-7 homolog A, crumbs cell polarity complex component , LIN-7A , LIN7 , MALS-1 , |

| MALS1 | . TIP-33 | . VELI1 |
|-------|----------|---------|
|       |          |         |

Background: Plays a role in establishing and maintaining the asymmetric distribution of channels and receptors at the plasma membrane of polarized cells. Forms membrane-associated multiprotein complexes that may regulate delivery and recycling of proteins to the correct membrane domains. The tripartite complex composed of LIN7 (LIN7A, LIN7B or LIN7C), CASK and APBA1 may have the potential to couple synaptic vesicle exocytosis to cell adhesion in brain. Ensures the proper localization of GRIN2B (subunit 2B of the NMDA receptor) to neuronal postsynaptic density and may function in localizing synaptic vesicles at synapses where it is recruited by beta-catenin and cadherin. Required to localize Kir2 channels, GABA transporter (SLC6A12) and EGFR/ERBB1, ERBB2, ERBB3 and ERBB4 to the basolateral membrane of epithelial cells.

| Molecular Weight: | 26 kDa            |
|-------------------|-------------------|
| Gene ID:          | 8825              |
| UniProt:          | 014910            |
| Pathways:         | Synaptic Membrane |

#### **Application Details**

| Application Notes: | WB: 1:1000-1:10000. ICC/IF: 1:100-1:1000. IHC-P: 1:100-1:1000. Optimal               |  |
|--------------------|--|--|
|                    | dilutions/concentrations should be determined by the researcher. Not tested in other |  |
|                    | applications.  |  |
| Comment:           | Positive Control: MCF-7  |  |
| Restrictions:      | For Research Use only  |  |

#### Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 mg/mL  |
| Buffer:            | 1XPBS (pH 7), 1 % BSA, 20 % Glycerol, 0.01 % Thimerosal  |
| Preservative:      | Thimerosal (Merthiolate)   |
| Precaution of Use: | This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage:           | 4 °C,-20 °C  |

### Handling

Storage Comment:

Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.