

## Datasheet for ABIN7468905

## anti-RPLP0 antibody



_					
	1//	r	Vİ	$\triangle$	۸/
	V		VI		/ V

Overview	
Quantity:	100 μL
Target:	RPLP0
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RPLP0 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	Recombinant protein encompassing a sequence within the center region of human RPLP0. The exact sequence is proprietary.
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by antigen-affinity chromatography.
Target Details	
Target:	RPLP0
Alternative Name:	ribosomal protein lateral stalk subunit P0 (RPLP0 Products)
Background:	Synonyms: ribosomal protein lateral stalk subunit P0 , L10E , LP0 , P0 , PRLP0 , RPP0
	Background: Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and
	The second secon

approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a
component of the 60S subunit. The protein, which is the functional equivalent of the E. coli L10
ribosomal protein, belongs to the L10P family of ribosomal proteins. It is a neutral
phosphoprotein with a C-terminal end that is nearly identical to the C-terminal ends of the acidic
ribosomal phosphoproteins P1 and P2. The P0 protein can interact with P1 and P2 to form a
pentameric complex consisting of P1 and P2 dimers, and a P0 monomer. The protein is located
in the cytoplasm. Transcript variants derived from alternative splicing exist, they encode the
same protein. As is typical for genes encoding ribosomal proteins, there are multiple processed
pseudogenes of this gene dispersed through the genome. [provided by RefSeq]

Molecular Weight:	34 kDa
Gene ID:	6175
UniProt <sup>.</sup>	P05388

## **Application Details**

Application Notes:	WB: 1:500-1:3000. IHC-P: 1:100-1:1000. Optimal dilutions/concentrations should be determined	
	by the researcher. Not tested in other applications.	
Comment:	Positive Control: 293T , A431	
Restrictions:	For Research Use only	

## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	1XPBS (pH 7), 20 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid

multiple freeze-thaw cycles.