antibodies -online.com





Datasheet for ABIN7468966 anti-SH3GLB2 antibody

| | i⊢₩ |
|--|-----|
| | |
| | |

| Overview | | |
|-------------------|----------------------------------------------------------------------------------------|--|
| Quantity: | 100 μL | |
| Target: | SH3GLB2 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This SH3GLB2 antibody is un-conjugated | |
| Application: | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) | |
| Product Details | | |
| Immunogen: | Recombinant protein encompassing a sequence within the center region of human SH3GLB2. | |
| | The exact sequence is proprietary. | |
| Isotype: | IgG | |
| Cross-Reactivity: | Human | |
| Purification: | Purified by antigen-affinity chromatography. | |
| Target Details | | |
| Target: | SH3GLB2 | |
| Alternative Name: | SH3 domain containing GRB2 like, endophilin B2 (SH3GLB2 Products) | |
| Background: | SH3 domain containing GRB2 like, endophilin B2 , PP6569 , PP9455 , RRIG1 | |
| Molecular Weight: | 44 kDa | |
| | | |

Target Details

| Gene ID: | 56904 | |
|---------------------|--------------------------------------------------------------------------------------------------|--|
| UniProt: | Q9NR46 | |
| Application Details | | |
| | | |
| Application Notes: | WB: 1:500-1:3000. IHC-P: 1:100-1:1000. Optimal dilutions/concentrations should be determined | |
| | by the researcher. Not tested in other applications. | |
| Comment: | Positive Control: NT2D1 , IMR32 , MCF-7 | |
| Restrictions: | For Research Use only | |
| | | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 0.98 mg/mL | |
| Buffer: | 1XPBS (pH 7), 1 % BSA, 20 % Glycerol, 0.01 % Thimerosal | |
| Preservative: | Thimerosal (Merthiolate) | |
| Precaution of Use: | This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE | |
| | which should be handled by trained staff only. | |
| | | |
| Storage: | 4 °C,-20 °C | |
| Storage Comment: | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage | |
| | (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid | |
| | multiple freeze-thaw cycles. | |
| | multiple meeze than cycles. | |