antibodies -online.com





Datasheet for ABIN7469292

anti-MBD4 antibody



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Overview		
Quantity:	100 μL	
Target:	MBD4	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC)	
Product Details		
Immunogen:	Recombinant protein encompassing a sequence within the center region of human MBD4. The exact sequence is proprietary.	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by antigen-affinity chromatography.	
Target Details		
Target:	MBD4	
Alternative Name:	MBD4 (MBD4 Products)	
Background:	Synonyms: methyl-CpG binding domain 4, DNA glycosylase , MED1	
	Background: DNA methylation is the major modification of eukaryotic genomes and plays an	
	essential role in mammalian development. Human proteins MECP2, MBD1, MBD2, MBD3, and	
	MBD4 comprise a family of nuclear proteins related by the presence in each of a methyl-CpG	

binding domain (MBD). Each of these proteins, with the exception of MBD3, is capable of		
binding specifically to methylated DNA. MBD4 may function to mediate the biological		
consequences of the methylation signal. In addition, MBD4 has protein sequence similarity to		
bacterial DNA repair enzymes and thus may have some function in DNA repair. Further, MBD4		
gene mutations are detected in tumors with primary microsatellite-instability (MSI), a form of		
genomic instability associated with defective DNA mismatch repair, and MBD4 gene meets 4 of		
5 criteria of a bona fide MIS target gene. [provided by RefSeq]		

Molecular Weight:	66 kDa
Gene ID:	8930
UniProt:	095243
Pathways:	DNA Damage Repair

Application Details

Application Notes:	WB: 1:1000-1:10000. ICC/IF: 1:100-1:1000. Optimal dilutions/concentrations should be
	determined by the researcher. Not tested in other applications.
Comment:	Positive Control: 3xFlag-human MBD4-transfected 293T cells
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	1XPBS (pH 7), 20 % Glycerol, 0.025 % ProClin 300
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.