

Datasheet for ABIN7469299

anti-COX15 antibody



Overview

100 μL
COX15
Human
Rabbit
Polyclonal
This COX15 antibody is un-conjugated
Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Recombinant protein encompassing a sequence within the center region of human COX15. The exact sequence is proprietary.
IgG
Human
Purified by antigen-affinity chromatography.
COX15
COX15 (COX15 Products)
Synonyms: cytochrome c oxidase assembly homolog COX15, CEMCOX2 Background: Cytochrome c oxidase (COX), the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. This

component is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes a protein which is not a structural subunit, but may be essential for the biogenesis of COX formation and may function in the hydroxylation of heme O, according to the yeast mutant studies. This protein is predicted to contain 5 transmembrane domains localized in the mitochondrial inner membrane. Alternative splicing of this gene generates two transcript variants diverging in the 3' region. [provided by RefSeq]

Molecular Weight:	46 kDa
Gene ID:	1355
UniProt:	Q7KZN9

Application Details

Application Notes:	WB: 1:500-1:3000. IHC-P: 1:100-1:1000. Optimal dilutions/concentrations should be determined
	by the researcher. Not tested in other applications.
Comment:	Positive Control: Jurkat
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	0.8 mg/mL
Buffer:	0.1M Tris-Glycine (pH 7), 20 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid

multiple freeze-thaw cycles.