## .-online.com antibodies

# Datasheet for ABIN7469457 anti-KIR2DS2 antibody



Overview

Quantity:	100 µL
Target:	KIR2DS2
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KIR2DS2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunofluorescence (IF)

#### Product Details

Immunogen:	Recombinant protein encompassing a sequence within the center region of human KIR2DS2. The exact sequence is proprietary.
lsotype:	lgG
Cross-Reactivity:	Human
Purification:	Purified by antigen-affinity chromatography.

#### Target Details

Target:	KIR2DS2
Alternative Name:	KIR2DS2 (KIR2DS2 Products)
Background:	Synonyms: killer cell immunoglobulin like receptor, two lg domains and short cytoplasmic tail 2 , 183Actl , CD158J , CD158b , KIR-2DS2 , NKAT-5 , NKAT5 , cl-49

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7469457 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

Molecular Weight:	34 kDa
	an important role in regulation of the immune response. [provided by RefSeq]
	several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to play
	TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for
	proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the
	signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR
	cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory
	immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S)
	KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular
	haplotypes, although several "framework" genes are found in all haplotypes (KIR3DL3, KIR3DP1,
	leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among
	highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb
	expressed by natural killer cells and subsets of T cells. The KIR genes are polymorphic and
	Background: Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins

merecalar mergina	
Gene ID:	100132285
UniProt:	P43631

#### Application Details

Application Notes:	WB: 1:500-1:3000. ICC/IF: 1:100-1:1000. IHC-P: 1:100-1:1000. Optimal dilutions/concentrations
	should be determined by the researcher. Not tested in other applications.
Comment:	Positive Control: Jurkat
Restrictions:	For Research Use only

### Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.1M Tris-Glycine ( pH 7), 20 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN7469457 | 09/09/2023 | Copyright antibodies-online. All rights reserved. Storage Comment:

Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.