## antibodies -online.com





Datasheet for ABIN7469613

## anti-SS18L1 antibody



Go to Product page

$\sim$					
	)\/e	r	٦\/	10	1///

Overview		
Quantity:	100 μL	
Target:	SS18L1	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SS18L1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC)	
Product Details		
Immunogen:	Recombinant protein encompassing a sequence within the center region of human CREST. The exact sequence is proprietary.	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by antigen-affinity chromatography.	
Target Details		
Target:	SS18L1	
Alternative Name:	CREST (SS18L1 Products)	
Background:	Synonyms: SS18L1 subunit of BAF chromatin remodeling complex, CREST, LP2261  Background: Synovial sarcomas occur most frequently in the extremities around large joints.  More than 90 % of cases have a recurrent and specific chromosomal translocation,	

Target Details	
	t(X,18)(p11.2,q11.2), in which the 5-prime end of the SS18 gene (MIM 600192) is fused in-frame to the 3-prime end of the SSX1 (MIM 312820), SSX2 (MIM 300192), or SSX4 (MIM 300326) gene. The SS18L1 gene is homologous to SS18.[supplied by OMIM]
Molecular Weight:	43 kDa
Gene ID:	26039
UniProt:	075177
Application Details	
Application Notes:	WB: 1:500-1:3000. ICC/IF: 1:100-1:1000. Optimal dilutions/concentrations should be determined by the researcher. Not tested in other applications.

Application Notes:	WB: 1:500-1:3000. ICC/IF: 1:100-1:1000. Optimal dilutions/concentrations should be determined	
	by the researcher. Not tested in other applications.	
Comment:	Positive Control: MCF-7, U87-MG NE nuclear extract, mouse brain, rat brain Validation: Orthogonal	
Restrictions:	For Research Use only	

## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	1XPBS ( pH 7), 1 % BSA, 20 % Glycerol, 0.01 % Thimerosal
Preservative:	Thimerosal (Merthiolate)
Precaution of Use:	This product contains Thimerosal (Merthiolate): a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.