antibodies.com

Datasheet for ABIN7477501 Recombinant anti-Inhibin alpha antibody (AA 233-362)




3 Images



1 mL
Inhibin alpha (INHA)
AA 233-362
Human
Rabbit
Recombinant Antibody
Monoclonal
This Inhibin alpha antibody is un-conjugated
Immunohistochemistry (IHC)
Recombinant fragment of human Inhibin alpha protein (around aa 233-362) (exact sequence is
proprietary)
MSVA-561R
lgG
Inhibin alpha (INHA)
Inhibin alpha (INHA) Inhibin Alpha (INHA Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7477501 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

-	
	validated for immunohistochemistry on 76 different Normal Tissues
UniProt:	P05111
Pathways:	Peptide Hormone Metabolism, Hormone Activity, Negative Regulation of Hormone Secretion
Application Details	
Application Notes:	IHC 1:75-1:150
Comment:	Positive Control: Adrenal gland: A moderate to strong INHA positivity should be seen in adrenocortical cells. Negative Control: Colon: INHA immunostaining should be completely absent.
Protocol:	Manual Protocol: Freshly cut sections should be used (less than 10 days between cutting and staining). Heat-induced antigen retrieval for 5 minutes in an autoclave at 121 °C in pH 7,8 Target Retrieval Solution buffer. Apply the antibody at a dilution of 1:50 at 37 °C for 60 minutes Visualization of bound antibody by the EnVision Kit (Dako, Agilent) according to the manufacturer's directions.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non- hazardous.



Immunohistochemistry

Image 1. The strongest inhibin alpha immunostaining among normal tissues is seen in the corpus luteum of the ovary

Immunohistochemistry

Image 2. Strong inhibin alpha immunostaining in all cells of an adrenocortical adenoma

Immunohistochemistry

Image 3. A strong inhibin alpha immunostaining is seen in the cyto and syncytiotrophoblast of the placenta of the first trimenon

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7477501 | 09/09/2023 | Copyright antibodies-online. All rights reserved.