antibodies -online.com





anti-GP120 antibody (Peroxidase (POD))



| \sim | | | | | |
|--------|----|----|----|-------------|---|
| () | VE | ۲۱ | /1 | \triangle | Λ |

| Quantity: | 1 mL | |
|---|---|--|
| Target: | GP120 (HIV-1 gp120) | |
| Reactivity: | Human Immunodeficiency Virus (HIV) | |
| Host: | Goat | |
| Clonality: | Polyclonal | |
| Conjugate: | This GP120 antibody is conjugated to Peroxidase (POD) | |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Enzyme Immunoassay (EIA) | |
| Product Details | | |
| Immunogen: | purified native gp120 from strain IIIB | |
| iriiridilogen. | purmed harive gp 120 from strain filb | |
| Isotype: | IgG | |
| | | |
| Isotype: | IgG | |
| Isotype: Specificity: | IgG Gp120 (predominate gp120 band in WB) | |
| Isotype: Specificity: | IgG Gp120 (predominate gp120 band in WB) Antiserum does not cross-react with human T or B cells UNINFECTED CELL REACTIVITY: | |
| Isotype: Specificity: Cross-Reactivity (Details): | IgG Gp120 (predominate gp120 band in WB) Antiserum does not cross-react with human T or B cells UNINFECTED CELL REACTIVITY: Antiserum is not reactive with human or bovine sera | |
| Isotype: Specificity: Cross-Reactivity (Details): | IgG Gp120 (predominate gp120 band in WB) Antiserum does not cross-react with human T or B cells UNINFECTED CELL REACTIVITY: Antiserum is not reactive with human or bovine sera This product consists of purified IgG fraction of the above antiserum covalently coupled to a | |
| Isotype: Specificity: Cross-Reactivity (Details): | IgG Gp120 (predominate gp120 band in WB) Antiserum does not cross-react with human T or B cells UNINFECTED CELL REACTIVITY: Antiserum is not reactive with human or bovine sera This product consists of purified IgG fraction of the above antiserum covalently coupled to a highly purified preparation of horseradish peroxidase (RZ >3) by methods. Care is taken to | |

substitution is 2-3.

Target Details

| Target: | GP120 (HIV-1 gp120) | |
|---------------------|---|--|
| Alternative Name: | Human Immunodeficiency Virus 1 Surface Glycoprotein (HIV-1 gp120 Products) | |
| Application Details | | |
| Application Notes: | TITER : > 1:1,000 by western blot immunoassay, This product is suitable for | |
| | immunohistochemical, immunoblotting and EIA applications. Working strength must be | |
| | determined by the user but a starting range of 1:20 - 1:200 is recommended for | |
| | immunochemistry and blotting while 1:200 - 1:1,000 is recommended for enzyme | |
| | immunoassays. | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | Lot specific | |
| Buffer: | This preparation is supplied in a phosphate saline buffer stabilized with bovine serum albumin | |
| | (10 mg/mL) and preserved with 0.002 % thimerosal. NOTE: sodium azide is a potent inhibitor of peroxidase! | |
| Preservative: | Thimerosal (Merthiolate) | |
| Precaution of Use: | This product contains Thimerosal (Merthiolate) and Sodium azide: POISONOUS AND | |
| | HAZARDOUS SUBSTANCES which should be handled by trained staff only. | |
| Storage: | 4 °C,-20 °C | |
| Storage Comment: | Recommended short term (<6 months) storage is liquid at 2-8°C. For longer term storage, aliquot and freeze. | |