antibodies .- online.com







anti-IL17C antibody (AA 101-197) (Cy5)



Overview

Quantity:	100 μL
Target:	IL17C
Binding Specificity:	AA 101-197
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IL17C antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human IL-17C
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	IL17C
Alternative Name:	IL-17C (IL17C Products)

Target Details

Background:	Synonyms: CX2, IL-17C, Interleukin-17C, Cytokine CX2, IL17C, UNQ561/PR01122
	Background: Cytokine that plays a crucial role in innate immunity of the epithelium, including to
	intestinal bacterial pathogens, in an autocrine manner. Stimulates the production of
	antibacterial peptides and proinflammatory molecules for host defense by signaling through
	the NF-kappa-B and MAPK pathways. Acts synergically with IL22 in inducing the expression of
	antibacterial peptides, including S100A8, S100A9, REG3A and REG3G. Synergy is also observed
	with TNF and IL1B in inducing DEFB2 from keratinocytes. Depending on the type of insult, may
	have both protective and pathogenic properties, either by maintaining epithelial homeostasis
	after an inflammatory challenge or by promoting inflammatory phenotype. Enhanced
	IL17C/IL17RE signaling may also lead to greater susceptibility to autoimmune diseases.
Gene ID:	27189
UniProt:	Q9P0M4
Pathways:	Cellular Response to Molecule of Bacterial Origin
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months