# antibodies -online.com





# anti-LCK antibody (AA 421-509)

3 Images



Go to Product page

$\sim$							
	1//	$\Box$	$r \setminus$	/ [	$\bigcirc$	1	٨,

Background:

Quantity:	100 μL
Target:	LCK
Binding Specificity:	AA 421-509
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LCK antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Flow Cytometry (FACS)
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human Lck
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Horse
Purification:	Purified by Protein A.
Target Details	
Target:	LCK
Alternative Name:	

Synonyms: LSK, YT16, IMD22, p56lck, pp58lck, Tyrosine-protein kinase Lck, Leukocyte C-

terminal Src kinase, Lymphocyte cell-specific protein-tyrosine kinase, Protein YT16, Protooncogene Lck, T cell-specific protein-tyrosine kinase, p56-LCK, LCK

Background: Non-receptor tyrosine-protein kinase that plays an essential role in the selection and maturation of developing T-cells in the thymus and in the function of mature T-cells. Plays a key role in T-cell antigen receptor (TCR)-linked signal transduction pathways. Constitutively associated with the cytoplasmic portions of the CD4 and CD8 surface receptors. Association of the TCR with a peptide antigen-bound MHC complex facilitates the interaction of CD4 and CD8 with MHC class II and class I molecules, respectively, thereby recruiting the associated LCK protein to the vicinity of the TCR/CD3 complex. LCK then phosphorylates tyrosines residues within the immunoreceptor tyrosine-based activation motifs (ITAM) of the cytoplasmic tails of the TCR-gamma chains and CD3 subunits, initiating the TCR/CD3 signaling pathway. Once stimulated, the TCR recruits the tyrosine kinase ZAP7, that becomes phosphorylated and activated by LCK. Following this, a large number of signaling molecules are recruited, ultimately leading to lymphokine production. LCK also contributes to signaling by other receptor molecules. Associates directly with the cytoplasmic tail of CD2, which leads to hyperphosphorylation and activation of LCK. Also plays a role in the IL2 receptor-linked signaling pathway that controls the T-cell proliferative response. Binding of IL2 to its receptor results in increased activity of LCK. Is expressed at all stages of thymocyte development and is required for the regulation of maturation events that are governed by both pre-TCR and mature alpha beta TCR. Phosphorylates other substrates including RUNX3, PTK2B/PYK2, the microtubule-associated protein MAPT, RHOH or TYROBP.

Gene ID: 3932

UniProt: P06239

Pathways: TCR Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin

Signaling Pathway, Transition Metal Ion Homeostasis, Positive Regulation of Endopeptidase

Activity, CXCR4-mediated Signaling Events, Thromboxane A2 Receptor Signaling

# **Application Details**

Application Notes: WB 1:300-5000

ELISA 1:500-1000


FCM 1:20-100

Restrictions: For Research Use only

# Handling


Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months	

# **Images**




# **Immunohistochemistry**

**Image 1.** Formalin-fixed and paraffin embedded human lung carcinoma labeled with Anti-Lck/p56-LCK Polyclonal Antibody, Unconjugated (ABIN748553) at 1:200 followed by conjugation to the secondary antibody and DAB staining.



### **ELISA**

**Image 2.** Antigen: 0.2  $\mu$ g/100  $\mu$ L Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000; Secondary: HRP conjugated Goat-Anti-Rabbit IgG at 1: 5000; TMB staining; Read the data in MicroplateReader by 450



# **ELISA**

Image 3. Antigen: 0.2ug/100ul, Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000, Secondary: HRP conjugated Goat-Anti-Rabbit IgG at 1: 5000, TMB staining, Read the data in MicroplateReader by 450nm