# antibodies - online.com







# anti-SPHK1 antibody (AA 1-100)





**Publications** 



0	1 /	-	K	/1	-	1 A
u	\/	$\vdash$	I \	/ I	$\vdash$	1/1

Quantity:	100 μL
Target:	SPHK1
Binding Specificity:	AA 1-100
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SPHK1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)),
	Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence
	(Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human SPHK1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Rabbit
Purification:	Purified by Protein A.

### **Target Details**

|--|

#### **Target Details**

Expiry Date:

l arget Details			
Alternative Name:	SPHK1 (SPHK1 Products)		
Background:	Synonyms: SPHK, Sphingosine kinase 1, SK 1, SPK 1, SPHK1, SPK		
	Background: Catalyzes the phosphorylation of sphingosine to form sphingosine 1-phosphate		
	(SPP), a lipid mediator with both intra- and extracellular functions. Also acts on D-erythro-		
	sphingosine and to a lesser extent sphinganine, but not other lipids, such as D,L-threo-		
	dihydrosphingosine, N,N-dimethylsphingosine, diacylglycerol, ceramide, or phosphatidylinositol.		
Gene ID:	8877		
UniProt:	Q9NYA1		
Pathways:	VEGF Signaling		
Application Details			
Application Notes:	WB 1:300-5000		
	ELISA 1:500-1000		
	IHC-P 1:200-400		
	IHC-F 1:100-500		
	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.		
Storage:	4 °C,-20 °C		
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		

12 months


Product cited in:

Tran, Jersmann, Truong, Hamon, Roscioli, Ween, Pitman, Pitson, Hodge, Reynolds, Hodge: "Disrupted epithelial/macrophage crosstalk via Spinster homologue 2-mediated S1P signaling may drive defective macrophage phagocytic function in COPD." in: **PLoS ONE**, Vol. 12, Issue 11, pp. e0179577, (2017) (PubMed).

Tani, Tabuchi, Hara: "Hair Cell Loss Induced by Sphingosine and a Sphingosine Kinase Inhibitor in the Rat Cochlea." in: **Neurotoxicity research**, Vol. 29, Issue 1, pp. 35-46, (2016) (PubMed).

Tran, Barnawi, Ween, Hamon, Roscioli, Hodge, Reynolds, Pitson, Davies, Haberberger, Hodge: "Cigarette smoke inhibits efferocytosis via deregulation of sphingosine kinase signaling: reversal with exogenous S1P and the S1P analogue FTY720." in: **Journal of leukocyte biology**, Vol. 100, Issue 1, pp. 195-202, (2016) (PubMed).

#### **Images**



#### **Western Blotting**

**Image 1.** Mouse heart lysates probed with SPHK1 Polyclonal Antibody, unconjugated at 1:30 overnight at 4°C followed by a conjugated secondary antibody at 1:10000 for 90 minutes at 37°C.