

Datasheet for ABIN7488952 **ENPP3 Protein (Fc Tag)**



Go to Product page

Overview

Quantity:	100 µg
Target:	ENPP3
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This ENPP3 protein is labelled with Fc Tag.

Product Details

Purpose:	Human ENPP3 / CD203c Protein, Fc Tag (active enzyme, MALS verified)
Sequence:	Leu 48 - Ile 875
Characteristics:	Human ENPP3 Protein, Fc Tag is expressed from human 293 cells (HEK293). It contains AA Leu 48 - Ile 875 (Accession # 014638-1).
Purity:	90 %
Endotoxin Level:	1.0 EU per µg
Grade:	MALS verified
Target Details	
Target:	ENPP3

Target.	
Alternative Name:	ENPP3 (ENPP3 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7488952 | 07/24/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Background:	The human NPP family contains seven members which can be classified into two groups
	according to their substrate preferences. The first class comprises the nucleotide-degrading
	proteins NPP1, 3 and 4. NPP3 (CD203c, ENPP3) is expressed in multiple organs, including on
	epithelial and mucosal surfaces, and notably on basophils and mast cells. Activation of
	basophils by antigen-bound IgE leads to release of inflammatory mediators and rapid
	upregulation of NPP3 to the cell surface. This protein is in fact a common marker for
	diagnosing allergen sensitivity with patient basophils by flow cytometry. Basophils and mast
	cells mediate the response to certain pathogens, as well as acute and chronic allergic
	reactions. Following activation, these cells release ATP, which further stimulates them in an
	autocrine manner. NPP3 upregulation serves to degrade ATP and suppress chronic allergic
	inflammation (but not the acute response).
Molecular Weight:	121.5 kDa
Pathways:	Regulation of Muscle Cell Differentiation, Negative Regulation of Transporter Activity
Application Details	
Comment:	This protein carries a human IgG1 Fc tag at the N-terminus. (Fc) The protein has a calculated
	MW of 121.5 kDa. The protein migrates as 150-190 kDa under reducing (R) condition (SDS-
	PAGE) due to glycosylation.
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Buffer:	50 mM Tris, 150 mM NaCl, pH 7.5
Storage:	-20 °C
Storage Comment:	-20°C