antibodies -online.com





CLEC4C Protein (AA 48-212) (Fc Tag)



Overview

Quantity:	500 μg
Target:	CLEC4C
Protein Characteristics:	AA 48-212
Origin:	Cynomolgus
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CLEC4C protein is labelled with Fc Tag.

Product Details

Cynomolgus BDCA-2 Protein
Tyr48-lle212
Recombinant Cynomolgus BDCA-2 Protein is expressed from HEK293 with hFc tag at the N-Terminus.It contains Tyr48-Ile212.
> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
0.22 µm filtered
Less than 1EU per µg by the LAL method.
Immobilized Cynomolgus BDCA-2, hFc Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Anti-BDCA-2 Antibody, hFc Tag with the EC50 of 13.1ng/ml determined by ELISA. See testing image for detail.

Target Details

Target:	CLEC4C
Alternative Name:	BDCA-2 (CLEC4C Products)
Background:	BDCA-2, BDCA-3, and BDCA-4. In blood, BDCA-2 and BDCA-4 are expressed on CD11c(-) CD123(bright) plasmacytoid dendritic cells, whereas BDCA-3 is expressed on small population of CD11c() CD123(-) dendritic cells. All three Ags are not detectable on a third blood dendritic cell population, which is CD1c() CD11c(bright) CD123(dim), or on any other cells in blood.BDCA-2 is completely down-regulated on plasmacytoid CD11c(-) CD123(bright) dendritic cells, BDCA-2 is rapidly internalized at 37 degrees C after mAb labeling.
Molecular Weight:	44.99 kDa. Due to glycosylation, the protein migrates to 53-63 kDa based on Tris-Bis PAGE result.
UniProt:	A0A2K5UWP4
Application Details	

Restrictions: For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from $0.22\mu m$ filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt.,-80°C for 3-6 months after reconstitution.,2-8°C for 2-7 days after reconstitution.,Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months