

Datasheet for ABIN7490759

HCST Protein (AA 19-48) (Fc Tag)





Go to Product page

| \sim | | | | |
|--------|----------------|-------|--------|----|
| () | Ive | r\ / | \cap | Λ. |
| \cup | $\lor \subset$ | I V I | \Box | ٧V |

| Quantity: | 100 μg |
|-------------------------------|--------------------------------------------|
| Target: | HCST |
| Protein Characteristics: | AA 19-48 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This HCST protein is labelled with Fc Tag. |

Product Details

| Purpose: | Recombinant human DAP10 protein with C-terminal human Fc tag | |
|------------------|------------------------------------------------------------------------------------------------------|--|
| Specificity: | DAP10 (Gln19-Pro48) hFc (Glu99-Ala330) | |
| Characteristics: | Extracellular Domain Protein | |
| Purification: | Purified from cell culture supernatant by affinity chromatography | |
| Purity: | The purity of the protein is greater than 95 % as determined by SDS-PAGE and Coomassie bluestaining. | |

Target Details

| Target: | HCST |
|--------------------------------------------------------------------------------------------|-----------------------|
| Alternative Name: | DAP10 (HCST Products) |
| Background: This gene encodes a transmembrane signaling adaptor that contains a YxxM motif | |

Storage:

Expiry Date:

Storage Comment:

| Target Details | |
|---------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| | cytoplasmic domain. The encoded protein may form part of the immune recognition receptor complex with the C-type lectin-like receptor NKG2D. As part of this receptor complex, this protein may activate phosphatidylinositol 3-kinase dependent signaling pathways through its intracytoplasmic YxxM motif. This receptor complex may have a role in cell survival and proliferation by activation of NK and T cell responses. Alternative splicing results in two transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008] |
| Molecular Weight: | predicted molecular mass of 29.1 kDa after removal of the signal peptide. The apparent molecular mass of DAP10-hFc is 25-55 kDa due to glycosylation. |
| UniProt: | Q9UBK5 |
| Application Details | |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Lyophilized |
| Buffer: | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose is added as protectants before lyophilization. |


Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for

use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

-20 °C,-80 °C

12 months



SDS-PAGE

Image 1. Human Protein, hFc Tag on SDS-PAGE under reducing condition.