antibodies .- online.com





Datasheet for ABIN7491157

CCR2 Protein (mFc Tag)



Overview

Quantity:	100 μg
Target:	CCR2
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CCR2 protein is labelled with mFc Tag.

Product Details

Purpose:	Recombinant human CCR2 protein with C-terminal mouse Fc tag
Specificity:	CCR2 (Met1-Ala42) mFc (Pro99-Lys330)
Characteristics:	Extracellular Domain Protein
Purity:	The purity of the protein is greater than 95 % as determined by SDS-PAGE and Coomassie blue staining.

Target Details

Target:	CCR2
Alternative Name:	CCR2 (CCR2 Products)
Background:	CC-CKR-2, MCP-1-R, CD192
	Description: The protein encoded by this gene is a receptor for monocyte chemoattractant
	protein-1, a chemokine which specifically mediates monocyte chemotaxis. Monocyte
	chemoattractant protein-1 is involved in monocyte infiltration in inflammatory diseases such as

Target Details

protein mediates agonist-dependent calcium mobilization and inhibition of adenylyl cyclase. This protein can also be a coreceptor with CD4 for HIV-1 infection. This gene is located in the chemokine receptor gene cluster region of chromosome 3. [provided by RefSeq, Aug 2017] Molecular Weight: predicted molecular mass of 31.1 kDa after removal of the signal peptide. The apparent molecular mass of CCR2-mFc is 35-55 kDa due to glycosylation. UniProt: P41597	Restrictions:	For Research Use only
protein mediates agonist-dependent calcium mobilization and inhibition of adenylyl cyclase. This protein can also be a coreceptor with CD4 for HIV-1 infection. This gene is located in the chemokine receptor gene cluster region of chromosome 3. [provided by RefSeq, Aug 2017] Molecular Weight: predicted molecular mass of 31.1 kDa after removal of the signal peptide. The apparent molecular mass of CCR2-mFc is 35-55 kDa due to glycosylation. UniProt: P41597 Pathways: cAMP Metabolic Process, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Leukocyte Mediated Immunity Positive Regulation P	Application Details	
protein mediates agonist-dependent calcium mobilization and inhibition of adenylyl cyclase. This protein can also be a coreceptor with CD4 for HIV-1 infection. This gene is located in the chemokine receptor gene cluster region of chromosome 3. [provided by RefSeq, Aug 2017] Molecular Weight: predicted molecular mass of 31.1 kDa after removal of the signal peptide. The apparent molecular mass of CCR2-mFc is 35-55 kDa due to glycosylation.	Pathways:	cAMP Metabolic Process, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process
protein mediates agonist-dependent calcium mobilization and inhibition of adenylyl cyclase. This protein can also be a coreceptor with CD4 for HIV-1 infection. This gene is located in the chemokine receptor gene cluster region of chromosome 3. [provided by RefSeq, Aug 2017] Molecular Weight: predicted molecular mass of 31.1 kDa after removal of the signal peptide. The apparent	UniProt:	P41597
protein mediates agonist-dependent calcium mobilization and inhibition of adenylyl cyclase. This protein can also be a coreceptor with CD4 for HIV-1 infection. This gene is located in the	Molecular Weight:	
rheumatoid arthritis as well as in the inflammatory response against tumors. The encoded		This protein can also be a coreceptor with CD4 for HIV-1 infection. This gene is located in the

Handling

Format:	Lyophilized
Buffer:	sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose is added as protectants before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Expiry Date:	12 months