antibodies -online.com





Datasheet for ABIN7491211

FURIN Protein (His tag)

Go to Product page

0	1 /	-	r.	/1	01	A /
	1//	\vdash	I \	/ I	-	\/\/

Quantity:	100 μg
Target:	FURIN
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This FURIN protein is labelled with His tag.

Product Details

Purpose:	Recombinant human FURIN protein with C-terminal 6xHis tag
Specificity:	FURIN (Gln27-Ala574) 6xHis tag
Characteristics:	Extracellular Domain Protein
Purity:	The purity of the protein is greater than 85 % as determined by SDS-PAGE and Coomassie blue staining.

Target Details

Target:	FURIN
Alternative Name:	FURIN (FURIN Products)
Background:	FUR, PACE, PCSK3, SPC1
	Description: This gene encodes a member of the subtilisin-like proprotein convertase family,
	which includes proteases that process protein and peptide precursors trafficking through
	regulated or constitutive branches of the secretory pathway. It encodes a type 1 membrane

bound protease that is expressed in many tissues, including neuroendocrine, liver, gut, and brain. The encoded protein undergoes an initial autocatalytic processing event in the ER and then sorts to the trans-Golgi network through endosomes where a second autocatalytic event takes place and the catalytic activity is acquired. Like other members of this convertase family, the product of this gene specifically cleaves substrates at single or paired basic residues. Some of its substrates include proparathyroid hormone, transforming growth factor beta 1 precursor, proalbumin, pro-beta-secretase, membrane type-1 matrix metalloproteinase, beta subunit of pro-nerve growth factor and von Willebrand factor. It is thought to be one of the proteases responsible for the activation of HIV envelope glycoproteins gp160 and gp140, and may play a role in tumor progression. Unlike SARS-CoV and other coronaviruses, the spike protein of SARS-CoV-2 is thought to be uniquely cleaved by this protease. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2020]

Molecular Weight:

predicted molecular mass of 61.0 kDa after removal of the signal peptide. The apparent molecular mass of FURIN-His is 55-70 kDa due to glycosylation.

UniProt:

P09958

Pathways:

Notch Signaling, Neurotrophin Signaling Pathway

Application Details

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Buffer:	sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose is added as protectants before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Expiry Date:	12 months