

### Datasheet for ABIN7491345

# Angiopoietin 2 Protein (ANGPT2) (AA 275-496) (His tag)



Image



| ( | ۱۱/ | e      | r\/ | Ì١ |               | ۱۸ |   |
|---|-----|--------|-----|----|---------------|----|---|
|   | , v | $\cup$ | V   | 1  | $\overline{}$ | V  | V |

| Quantity:                     | 100 μg  |
|-------------------------------|---|
| Target:                       | Angiopoietin 2 (ANGPT2)                               |
| Protein Characteristics:      | AA 275-496  |
| Origin:                       | Human   |
| Source:                       | HEK-293 Cells   |
| Protein Type:                 | Recombinant   |
| Purification tag / Conjugate: | This Angiopoietin 2 protein is labelled with His tag. |

#### **Product Details**

| Purpose:         | Recombinant Human ANGPT2(275-496) Protein with C-terminal 6xHis tag                                   |  |
|------------------|---|--|
| Specificity:     | ANGPT2 (Lys275-Phe496) 6xHis tag  |  |
| Characteristics: | Extracellular Domain Protein  |  |
| Purification:    | Purified from cell culture supernatant by affinity chromatography                                     |  |
| Purity:          | The purity of the protein is greater than 85 % as determined by SDS-PAGE and Coomassie blue staining. |  |

#### **Target Details**

| Target:           | Angiopoietin 2 (ANGPT2)   |
|-------------------|---|
| Alternative Name: | ANGPT2 (ANGPT2 Products)  |
| Background:       | This gene belongs to the angiopoietin family of growth factors. The protein encoded by this |

| gene is an antagonist of angiopoietin 1, and both angiopoietin 1 and angiopoietin 2 are ligands |  |  |
|---|--|--|
| for the endothelial TEK receptor tyrosine kinase. Angiopoietin 2 is upregulated in multiple     |  |  |
| inflammatory diseases and is implicated in the direct control of inflammation-related signaling |  |  |
| pathways. The encoded protein affects angiogenesis during embryogenesis and tumorigenesis,      |  |  |
| disrupts the vascular remodeling ability of angiopoietin 1, and may induce endothelial cell     |  |  |
| apoptosis. This gene serves a prognostic biomarker for acute respiratory distress syndrome.     |  |  |
| [provided by RefSeq, Aug 2020]  |  |  |

Molecular Weight:

predicted molecular mass of 26.3 kDa after removal of the signal peptide. The apparent molecular mass of ANGPT2(275-496)-His is 25-35 kDa due to glycosylation.

UniProt:

015123

Pathways:

**RTK Signaling** 


# **Application Details**

Restrictions:

For Research Use only

# Handling

| Format:          | Lyophilized  |
|------------------|--|
| Buffer:          | Lyophilized from sterile PBS, pH 7.4. Normally $5\% - 8\%$ trehalose is added as protectants before lyophilization.  |
| Storage:         | -20 °C,-80 °C  |
| Storage Comment: | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).  Lyophilized proteins are shipped at ambient temperature. |
| Expiry Date:     | 12 months  |



#### **SDS-PAGE**

**Image 1.** Human AN(275-496) Protein, His Tag on SDS-PAGE under reducing condition.