

[Go to Product page](#)

## Datasheet for ABIN7491621 **CXCR2 Protein**

### Overview

Quantity:	100 µg
Target:	CXCR2
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Synthetic

### Product Details

Purpose:	Human CXCR2 full length protein-synthetic nanodisc
Characteristics:	Full Length Transmembrane Proteins (synthetic Nanodisc)

### Target Details

Target:	CXCR2
Alternative Name:	CXCR2 ( <a href="#">CXCR2 Products</a> )
Background:	<p>CD182, CDw128b, CMKAR2, IL8R2, IL8RA, IL8RB</p> <p>Description: The protein encoded by this gene is a member of the G-protein-coupled receptor family. This protein is a receptor for interleukin 8 (IL8). It binds to IL8 with high affinity, and transduces the signal through a G-protein activated second messenger system. This receptor also binds to chemokine (C-X-C motif) ligand 1 (CXCL1/MGSA), a protein with melanoma growth stimulating activity, and has been shown to be a major component required for serum-dependent melanoma cell growth. This receptor mediates neutrophil migration to sites of inflammation. The angiogenic effects of IL8 in intestinal microvascular endothelial cells are found to be mediated by this receptor. Knockout studies in mice suggested that this receptor</p>

## Target Details

controls the positioning of oligodendrocyte precursors in developing spinal cord by arresting their migration. This gene, IL8RA, a gene encoding another high affinity IL8 receptor, as well as IL8RBP, a pseudogene of IL8RB, form a gene cluster in a region mapped to chromosome 2q33-q36. Alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq, Nov 2009]

Molecular Weight: The human full length CXCR2 protein has a MW of 40.8 kDa

UniProt: [P25025](#)

Pathways: [cAMP Metabolic Process](#)

## Application Details

Application Notes:

- Applications for VLPs:
- ELISA
- SPR affinity analysis
- Phage display screening
- Immunization
- Cell based assays
- CAR-T cell screening
- Protein crystal structure analysis

Comment: Synthetic Nanodisc can be prepared directly from the cells. The polymers used during this process have a dual function. It dissolves the cell membranes, like the detergent, and uses cellular phospholipids to form Nanodisc around the membrane proteins. The target protein embedded Nanodiscs can then be purified.

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Supplied in nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0)

Storage: -20 °C, -80 °C

Storage Comment: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).  
Lyophilized proteins are shipped at ambient temperature.

Expiry Date: 12 months