

[Go to Product page](#)

Datasheet for ABIN7491629 CXCR6 Protein

Overview

Quantity:	100 µg
Target:	CXCR6
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Synthetic

Product Details

Purpose:	Human CXCR6 full length protein-synthetic nanodisc
Characteristics:	Full Length Transmembrane Proteins (synthetic Nanodisc)

Target Details

Target:	CXCR6
Alternative Name:	CXCR6 (CXCR6 Products)
Background:	<p>BONZO, CD186, CDw186, STRL33, TYMSTR</p> <p>Description: The protein encoded by this gene is a G protein-coupled receptor with seven transmembrane domains that belongs to the CXC chemokine receptor family. This family also includes CXCR1, CXCR2, CXCR3, CXCR4, CXCR5, and CXCR7. This gene, which maps to the chemokine receptor gene cluster, is expressed in several T lymphocyte subsets and bone marrow stromal cells. The encoded protein and its exclusive ligand, chemokine ligand 16 (CCL16), are part of a signalling pathway that regulates T lymphocyte migration to various peripheral tissues (the liver, spleen red pulp, intestine, lungs, and skin) and promotes cell-cell interaction with dendritic cells and fibroblastic reticular cells. CXCR6/CCL16 also controls the</p>

Target Details

localization of resident memory T lymphocytes to different compartments of the lung and maintains airway resident memory T lymphocytes, which are an important first line of defense against respiratory pathogens. The encoded protein serves as an entry coreceptor used by HIV-1 and SIV to enter target cells, in conjunction with CD4. [provided by RefSeq, Aug 2020]

Molecular Weight: The human full length CXCR6 protein has a MW of 39.3 kDa

UniProt: [O00574](#)

Application Details

Application Notes:

- Applications for VLPs:
- ELISA
- SPR affinity analysis
- Phage display screening
- Immunization
- Cell based assays
- CAR-T cell screening
- Protein crystal structure analysis

Comment: Synthetic Nanodisc can be prepared directly from the cells. The polymers used during this process have a dual function. It dissolves the cell membranes, like the detergent, and uses cellular phospholipids to form Nanodisc around the membrane proteins. The target protein embedded Nanodiscs can then be purified.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Supplied in nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0)

Storage: -20 °C, -80 °C

Storage Comment: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).
Lyophilized proteins are shipped at ambient temperature.

Expiry Date: 12 months