

Datasheet for ABIN7491745

STEAP2 Protein**2** Images[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	STEAP2
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Synthetic Nanodisc

Product Details

Purpose:	Human STEAP2 full length protein-synthetic nanodisc
Characteristics:	Unlike other membrane scaffold protein (MSP) Nanodisc on the market, our synthetic Nanodisc can be prepared directly from the cells. The polymers used during this process have a dual function. It dissolves the cell membranes, like the detergent, and uses cellular phospholipids to form Nanodisc around the membrane proteins. The target protein embedded Nanodiscs can then be purified.

Target Details

Target:	STEAP2
Alternative Name:	STEAP2 (STEAP2 Products)
Background:	A member of the STEAP family and encodes a multi-pass membrane protein that localizes to the Golgi complex, the plasma membrane, and the vesicular tubular structures in the cytosol. A highly similar protein in mouse has both ferrireductase and cupric reductase activity, and stimulates the cellular uptake of both iron and copper in vitro. Increased transcriptional expression of the human gene is associated with prostate cancer progression. Alternate

Target Details


	transcriptional splice variants, encoding different isoforms, have been characterized.
Molecular Weight:	The human full length STEAP2 protein has a MW of 56.1 kDa
UniProt:	Q8NFT2
Pathways:	Transition Metal Ion Homeostasis

Application Details

Comment:	<p>Advantages of Synthetic Nanodiscs:</p> <ul style="list-style-type: none">• Highly purified membrane proteins• High solubility in aqueous solutions• High stability• Proteins are in a native membrane environment and remain biologically active• No detergent and can be used for cell-based assays• No MSP backbone proteins <p>Limitations of Synthetic Nanodiscs:</p> <ul style="list-style-type: none">• Intolerant to acids and high concentrations of divalent metal ions
Restrictions:	For Research Use only

Handling


Format:	Lyophilized
Buffer:	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5 % - 8 % trehalose is added as protectants before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Expiry Date:	12 months



SDS-PAGE

Image 1. Human ST-Nanodisc, Flag Tag on SDS-PAGE

ELISA assay to evaluate STEAP2-Nanodisc
0.2µg Human STEAP2-Nanodisc per well



ELISA

Image 2. Elisa plates were pre-coated with Flag Tag ST-Nanodisc (0.2 µg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with ST-Nanodisc is 9.198 ng/mL.