

Datasheet for ABIN750148

anti-CSF1R antibody (AA 201-300) (Cy5.5)



[Go to Product page](#)

1 Publication

Overview

Quantity:	100 µL
Target:	CSF1R
Binding Specificity:	AA 201-300
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CSF1R antibody is conjugated to Cy5.5
Application:	Flow Cytometry (FACS), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MCSF receptor
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat
Purification:	Purified by Protein A.

Target Details

Target:	CSF1R
Alternative Name:	MCSF Receptor (CSF1R Products)

Target Details

Background:	Synonyms: FMS, CSFR, FIM2, HDLS, C-FMS, CD115, CSF-1R, M-CSF-R, Macrophage colony-stimulating factor 1 receptor, CSF-1 receptor, CSF-1-R, Proto-oncogene c-Fms, CSF1R, CSF1R_HUMAN Background: The protein encoded by this gene is the receptor for colony stimulating factor 1, a cytokine which controls the production, differentiation, and function of macrophages. This receptor mediates most if not all of the biological effects of this cytokine. Ligand binding activates the receptor kinase through a process of oligomerization and transphosphorylation. The encoded protein is a tyrosine kinase transmembrane receptor and member of the CSF1/PDGF receptor family of tyrosine-protein kinases. Mutations in this gene have been associated with a predisposition to myeloid malignancy. The first intron of this gene contains a transcriptionally inactive ribosomal protein L7 processed pseudogene oriented in the opposite direction. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2013]
Gene ID:	1436
UniProt:	P07333
Pathways:	RTK Signaling , Inositol Metabolic Process , Cell-Cell Junction Organization

Application Details

Application Notes:	FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months

Publications

Product cited in: Gorden, Saha, Khammanivong, Schwartz, Dickerson: "Lysosomal drug sequestration as a mechanism of drug resistance in vascular sarcoma cells marked by high CSF-1R expression." in: **Vascular cell**, Vol. 6, pp. 20, (2014) ([PubMed](#)).