antibodies -online.com





Datasheet for ABIN7504143 anti-Follistatin antibody



Image



()	ve	K\ /		A .
	\cup	1 V/	Щ.	V۷

Overview		
Quantity:	100 μg	
Target:	Follistatin (FST)	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This Follistatin antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC)	
Product Details		
Immunogen:	Recognizes human Follistatin isoform FS288, raised against recombinant human FS288.	


Immunogen:	Recognizes human Follistatin isoform FS288, raised against recombinant human FS288.
Isotype:	lgG1
Specificity:	Follistatin is a single-chain glycosylated protein that inhibits follicle stimulating hormone (FSH)
	release. Alternative splicing of Follistatin mRNA yields two isoforms, FS315 and FS288. FS288
	is the main cell-surface form and binds to surface heparin sulphate proteoglycans. FST/4282
	recognizes recombinant human Follistatin 288, allowing for detection of FSH levels using
	various analysis methods. This antibody also works in a two site ELISA with FST/4281.
Cross-Reactivity (Details):	Human.
Purification:	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G.

Target Details

Target: Follistatin (FST)

Target Details	
Alternative Name:	FST (FST Products)
Background:	Follistatin, FS, FST, Active-binding protein, Follistatin / Activin Binding Protein Cellular localisation: Secreted.
Molecular Weight:	37kDa
Gene ID:	10468, 9914
UniProt:	P19883
Pathways:	Negative Regulation of Hormone Secretion
Application Details	
Application Notes:	Positive Control: Human testis or ovary. Known Application: ELISA (1 μ g/mL), Immunocytochemistry (70 μ g/mL), Western Blot (1-2 μ g/mL), Optimal dilution for a specific application should be determined.
Restrictions:	For Research Use only
Handling	

Concentration:	200 μg/mL
Buffer:	Prepared in 10 mM PBS with 0.05 % BSA and 0.05 % azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8 °C. Antibody is stable for 24 months. Non-hazardous. Also available WITHOUT BSA & azide at 1.0mg/ml.
Expiry Date:	24 months



Western Blotting

Image 1. SDS-PAGE Analysis Purified Follistatin Mouse Monoclonal Antibody (FST/4282). Confirmation of Purity and Integrity of Antibody.