# antibodies .- online.com







## CD37 Protein (AA 113-240) (His tag)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μg
Target:	CD37 (TSPAN26)
Protein Characteristics:	AA 113-240
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD37 protein is labelled with His tag.

#### **Product Details**

Purpose:	Human CD37 Protein	
Sequence:	Ala113-Asn240	
Characteristics:	Recombinant Human CD37 Protein is expressed from HEK293 with His tag at the C-Terminus.It contains Ala113-Asn240.	
Purity:	> 95 % as determined by Tris-Bis PAGE	
Sterility:	0.22 µm filtered	
Endotoxin Level:	Less than 1EU per μg by the LAL method.	

#### **Target Details**

Target:	CD37 (TSPAN26)
Alternative Name:	CD37 (TSPAN26 Products)

### **Target Details**

Expiry Date:

Currently, several monoclonal antibodies targeting CD37 as well as chimeric antigen recebased immunotherapies are being developed and investigated in clinical trials. Given the role of CD37 in the biology of B cells, it seems that CD37 constitutes more than a docking for monoclonal antibodies, and targeting this molecule may provide additional benefit to relapsed or refractory patients.  Molecular Weight: 15.72 kDa. Due to glycosylation, the protein migrates to 25-40 kDa based on Tris-Bis PAG result.  Pathways: Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response  Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/recommended. Dissolve the lyophilized protein in distilled water.  Buffer: Lyophilized from 0.22 µm filtered solution in PBS ( pH 7.4). Normally 8 % trehalose is add protectant before lyophilization.  Storage: -20 °C,-80 °C  Storage Comment: -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after	g =		
result.  Pathways: Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Im Response  Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/recommended. Dissolve the lyophilized protein in distilled water.  Buffer: Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is add protectant before lyophilization.  Storage: -20 °C,-80 °C  Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt., -80 °C for 3-6 months after reconstitution., 2-8 °C for 2-7 days after reconstitution., Recommend to aliquot the protein	Background:	molecular target exploited in the immunotherapy of B cell-derived lymphomas and leukemia. Currently, several monoclonal antibodies targeting CD37 as well as chimeric antigen receptor-based immunotherapies are being developed and investigated in clinical trials. Given the unique role of CD37 in the biology of B cells, it seems that CD37 constitutes more than a docking point for monoclonal antibodies, and targeting this molecule may provide additional benefit to	
Application Details  Restrictions: For Research Use only  Handling  Format: Lyophilized  Reconstitution: Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/recommended. Dissolve the lyophilized protein in distilled water.  Buffer: Lyophilized from 0.22 µm filtered solution in PBS ( pH 7.4). Normally 8 % trehalose is add protectant before lyophilization.  Storage: -20 °C,-80 °C  Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt., -80 °C for 3-6 months after reconstitution., 2-8 °C for 2-7 days after reconstitution., Recommend to aliquot the protein	Molecular Weight:	15.72 kDa. Due to glycosylation, the protein migrates to 25-40 kDa based on Tris-Bis PAGE result.	
Restrictions:       For Research Use only         Handling       Format:       Lyophilized         Reconstitution:       Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/recommended. Dissolve the lyophilized protein in distilled water.         Buffer:       Lyophilized from 0.22 μm filtered solution in PBS ( pH 7.4). Normally 8 % trehalose is add protectant before lyophilization.         Storage:       -20 °C, -80 °C         Storage Comment:       -20 to -80 °C for 12 months as supplied from date of receipt., -80 °C for 3-6 months after reconstitution., 2-8 °C for 2-7 days after reconstitution., Recommend to aliquot the protein	Pathways:	Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response	
Format:  Lyophilized  Reconstitution:  Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/recommended. Dissolve the lyophilized protein in distilled water.  Buffer:  Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is add protectant before lyophilization.  Storage:  -20 °C,-80 °C  Storage Comment:  -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein	Application Details		
Format:  Lyophilized  Reconstitution:  Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/recommended. Dissolve the lyophilized protein in distilled water.  Buffer:  Lyophilized from 0.22 µm filtered solution in PBS ( pH 7.4). Normally 8 % trehalose is add protectant before lyophilization.  Storage:  -20 °C,-80 °C  Storage Comment:  -20 to -80 °C for 12 months as supplied from date of receipt., -80 °C for 3-6 months after reconstitution., 2-8 °C for 2-7 days after reconstitution., Recommend to aliquot the protein	Restrictions:	For Research Use only	
Reconstitution:  Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/recommended. Dissolve the lyophilized protein in distilled water.  Buffer:  Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is add protectant before lyophilization.  Storage:  -20 °C,-80 °C  Storage Comment:  -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein	Handling		
recommended. Dissolve the lyophilized protein in distilled water.  Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is add protectant before lyophilization.  Storage:  -20 °C,-80 °C  Storage Comment:  -20 to -80 °C for 12 months as supplied from date of receipt., -80 °C for 3-6 months after reconstitution., 2-8 °C for 2-7 days after reconstitution., Recommend to aliquot the protein	Format:	Lyophilized	
protectant before lyophilization.  Storage: -20 °C,-80 °C  Storage Comment: -20 to -80 °C for 12 months as supplied from date of receipt., -80 °C for 3-6 months after reconstitution., 2-8 °C for 2-7 days after reconstitution., Recommend to aliquot the protein	Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu$ g/mL is recommended. Dissolve the lyophilized protein in distilled water.	
Storage Comment:  -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein	Buffer:	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.	
reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein	Storage:	-20 °C,-80 °C	
	Storage Comment:	reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into	

12 months