

Datasheet for ABIN7504274 CD47 Protein (CD47) (AA 19-139)



Overview

Quantity:	100 μg
Target:	CD47
Protein Characteristics:	AA 19-139
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant

Product Details

Purpose:	Human CD47 Protein
Sequence:	Gln19-Pro139
Characteristics:	Recombinant Human CD47 Protein is expressed from HEK293 without tag.lt contains Gln19-Pro139.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 μm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.

Target Details

Target:	CD47
Alternative Name:	CD47 (CD47 Products)
Background:	CD47 (Cluster of Differentiation 47) also known as integrin associated protein (IAP) is a

Storage:

Expiry Date:

Storage Comment:

-20 °C,-80 °C

12 months

Target Details	
	transmembrane protein that in humans is encoded by the CD47 gene. CD47 belongs to the immunoglobulin superfamily and partners with membrane integrins and also binds the ligands thrombospondin-1 (TSP-1) and signal-regulatory protein alpha (SIRPa).CD-47 acts as a don't eat me signal to macrophages of the immune system which has made it a potential therapeutic target in some cancers.
Molecular Weight:	13.72 kDa. Due to glycosylation, the protein migrates to 35-48 kDa based on Tris-Bis PAGE result.
UniProt:	Q08722-1
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.

-20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after

smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into