

## Datasheet for ABIN7504308

# DKK1 Protein (AA 32-142, N-Term) (mFc Tag)



#### Overview

| Quantity:                     | 100 μg                                      |
|-------------------------------|---|
| Target:                       | DKK1  |
| Protein Characteristics:      | AA 32-142, N-Term                           |
| Origin:                       | Human                                       |
| Source:                       | HEK-293 Cells                               |
| Protein Type:                 | Recombinant                                 |
| Purification tag / Conjugate: | This DKK1 protein is labelled with mFc Tag. |

### **Product Details**

| Purpose:         | Human DKK1 N terminal Domain Protein   |
|------------------|--|
| Sequence:        | Thr32-Asp142   |
| Characteristics: | Recombinant Human DKK1 N terminal Domain Protein is expressed from HEK293 with mFc tag at the C-Terminus.It contains Thr32-Asp142. |
| Purity:          | > 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC   |
| Sterility:       | 0.22 µm filtered   |
| Endotoxin Level: | Less than 1EU per μg by the LAL method.  |

## Target Details

| Target:           | DKK1                 |
|-------------------|----------------------|
| Alternative Name: | DKK1 (DKK1 Products) |

### **Target Details**

Expiry Date:

12 months

| Dickkopf-1 (Dkk1), the founding and best-studied member of the Dkk family, functions as an antagonist of canonical Wnt/β-catenin. Dkk1 is considered to play a broad role in a variety of biological processes. |
|---|
| 37.35 kDa. Due to glycosylation, the protein migrates to 45-50 kDa based on Tris-Bis PAGE result.   |
| 094907  |
| WNT Signaling, Regulation of Muscle Cell Differentiation, Positive Regulation of fat Cell Differentiation   |
|   |
| For Research Use only   |
|   |
| Lyophilized   |
| Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water.  |
| Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.  |
| -20 °C,-80 °C   |
| -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into                       |
|   |