# antibodies -online.com





Datasheet for ABIN7504423

## ICAM1 Protein (AA 25-476) (His tag)



#### Overview

Quantity:	100 μg
Target:	ICAM1
Protein Characteristics:	AA 25-476
Origin:	Dog
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ICAM1 protein is labelled with His tag.

#### **Product Details**

Purpose:	Canine ICAM-1/CD54 Protein
Sequence:	Gln25-Asn476
Characteristics:	Recombinant Canine ICAM-1/CD54 Protein is expressed from HEK293 with His tag at the C-Terminus.It contains Gln25-Asn476.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 µm filtered
Endotoxin Level:	Less than 1EU per µg by the LAL method.

### **Target Details**

Target:	ICAM1
Alternative Name:	ICAM-1 (ICAM1 Products)

#### **Target Details**

rarget Details	
Background:	Intercellular adhesion molecule-1 (ICAM-1, CD54) is a 90 kDa member of the immunoglobulin (Ig) superfamily and is critical for the firm arrest and transmigration of leukocytes out of blood vessels and into tissues. ICAM-1 is constitutively present on endothelial cells, but its expression is increased by proinflammatory cytokines. The endothelial expression of ICAM-1 is increased in atherosclerotic and transplant-associated atherosclerotic tissue and in animal models of atherosclerosis.
Molecular Weight:	51.08 kDa. Due to glycosylation, the protein migrates to 80-110 kDa based on Tris-Bis PAGE result.
UniProt:	P33729
Pathways:	Cellular Response to Molecule of Bacterial Origin, Regulation of Actin Filament Polymerization, Carbohydrate Homeostasis, Regulation of Leukocyte Mediated Immunity, Thromboxane A2 Receptor Signaling
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu$ g/mL is recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22 $\mu m$ filtered solution in PBS ( pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months