

## Datasheet for ABIN7504439 IL1F10 Protein (AA 1-152) (His tag)



| Overview                      |   |
|-------------------------------|---|
| Quantity:                     | 100 µg  |
| Target:                       | IL1F10  |
| Protein Characteristics:      | AA 1-152  |
| Origin:                       | Human   |
| Source:                       | Yeast   |
| Protein Type:                 | Recombinant   |
| Purification tag / Conjugate: | This IL1F10 protein is labelled with His tag.   |
| Product Details               |   |
| Purpose:                      | Human IL-1F10/IL-38 Protein   |
| Sequence:                     | Met1-Trp152   |
| Characteristics:              | Recombinant Human IL-1F10/IL-38 Protein is expressed from Yeast with His tag at the C-<br>terminus.It contains Met1-Trp152. |
| Purity:                       | > 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC  |
| Sterility:                    | 0.22 µm filtered  |
| Endotoxin Level:              | Less than 1EU per $\mu$ g by the LAL method.  |
| Target Details                |   |
| Target:                       | IL1F10  |
| Alternative Name:             | IL-1F10 (IL1F10 Products)   |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7504439 | 07/24/2024 | Copyright antibodies-online. All rights reserved.

| Background:         | Interleukin (IL)-38, a newly discovered IL-1 family cytokine, is expressed in several tissues and secreted by various cells. IL-38 has recently been reported to exert an anti-inflammatory function by binding to several receptors, including interleukin-36 receptor (IL-36R), interleukin-1 receptor accessory protein-like 1 (IL-1RAPL1), and interleukin-1 receptor 1 (IL-1R1) to block binding with other pro-inflammatory cytokines and inhibit subsequent signaling pathways, thereby regulating the differentiation and function of T cells, peripheral blood mononuclear cells, macrophages, and dendritic cells. |
|---------------------|--|
| Molecular Weight:   | 18.07 kDa. Due to glycosylation, the protein migrates to 20-23 kDa based on Tris-Bis PAGE result.  |
| Application Details |  |
| Restrictions:       | For Research Use only  |
| Handling            |  |
| Format:             | Lyophilized  |
| Reconstitution:     | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu$ g/mL is recommended. Dissolve the lyophilized protein in distilled water.  |
| Buffer:             | Lyophilized from 0.22 µm filtered solution in PBS ( pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization.  |
| Storage:            | -20 °C,-80 °C  |
| Storage Comment:    | -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.  |
| Expiry Date:        | 12 months  |