antibodies -online.com





PDGFRB Protein (AA 32-531) (His tag)



Overview

Quantity:	100 μg
Target:	PDGFRB
Protein Characteristics:	AA 32-531
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PDGFRB protein is labelled with His tag.

Product Details

Purpose:	Rat PDGF R beta/CD140b Protein
Sequence:	Leu32-Val531
Characteristics:	Recombinant Rat PDGF R beta/CD140b Protein is expressed from HEK293 with His tag at the C-terminus.It contains Leu32-Val531.
Purity:	> 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC
Sterility:	0.22 μm filtered
Endotoxin Level:	Less than 1EU per μg by the LAL method.

Target Details

Target:	PDGFRB
Alternative Name:	PDGF R beta (PDGFRB Products)

Target Details

Expiry Date:

. a. gat z atama	
Background:	PDGF is a major serum mitogen that can exist as a homo or hetero-dimeric protein consisting of disulfide-linked PDGF-A and PDGF-B chains. The PDGF-AA,PDGF-BB and PDGF-AB isoforms have been shown to bind to two distinct cell surface PDGF receptors with different affinities.
	Where as PDGF R alpha binds all three PDGF isoforms with high affinity, PDGF R beta binds
	PDGF-BB only with high-affinity. PDGF R plays an essential role in blood vessel development by
	promoting proliferation, migration and recruitment of pericytes and smooth muscle cells to
	endothelial cells. Plays a role in the migration of vascular smooth muscle cells and the
	formation of neointima at vascular injury sites.
Molecular Weight:	57.36 kDa. Due to glycosylation, the protein migrates to 80-110 kDa based on Tris-Bis PAGE result.
UniProt:	Q05030
Pathways:	Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling
	Pathway, Inositol Metabolic Process, Glycosaminoglycan Metabolic Process, Smooth Muscle
	Cell Migration, Platelet-derived growth Factor Receptor Signaling
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is
	recommended. Dissolve the lyophilized protein in distilled water.
Buffer:	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as
	protectant before lyophilization.
Storage:	-20 °C,-80 °C
Storage Comment:	-20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after
	reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into
	smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

12 months