

[Go to Product page](#)

Datasheet for ABIN7504541

TIE1 Protein (AA 22-759) (His tag)

Overview

| | |
|-------------------------------|---|
| Quantity: | 100 µg |
| Target: | TIE1 |
| Protein Characteristics: | AA 22-759 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This TIE1 protein is labelled with His tag. |

Product Details

| | |
|------------------|--|
| Purpose: | Human TIE1 Protein |
| Sequence: | Ala22-Gln759 |
| Characteristics: | Recombinant Human TIE1 Protein is expressed from HEK293 with His tag at the C-Terminus.It contains Ala22-Gln759. |
| Purity: | > 95 % as determined by Tris-Bis PAGE,> 95 % as determined by HPLC |
| Sterility: | 0.22 µm filtered |
| Endotoxin Level: | Less than 1EU per µg by the LAL method. |

Target Details

| | |
|-------------------|--|
| Target: | TIE1 |
| Alternative Name: | TIE1 (TIE1 Products) |

Target Details

| | |
|-------------------|--|
| Background: | TIE1 is a cell surface protein expressed in endothelial cells. Involved in angiogenesis and lymphangiogenesis, including morphogenesis of lymphatic valves, TIE1 is important for lymphatic system functional integrity. |
| Molecular Weight: | 80.93 kDa. Due to glycosylation, the protein migrates to 81-100 kDa based on Tris-Bis PAGE result. |
| Pathways: | RTK Signaling |

Application Details

| | |
|---------------|-----------------------|
| Restrictions: | For Research Use only |
|---------------|-----------------------|

Handling

| | |
|------------------|---|
| Format: | Lyophilized |
| Reconstitution: | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/mL is recommended. Dissolve the lyophilized protein in distilled water. |
| Buffer: | Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8 % trehalose is added as protectant before lyophilization. |
| Storage: | -20 °C,-80 °C |
| Storage Comment: | -20 to -80°C for 12 months as supplied from date of receipt., -80°C for 3-6 months after reconstitution., 2-8°C for 2-7 days after reconstitution., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |
| Expiry Date: | 12 months |