
# antibodies -online.com





# beta-2 Microglobulin Protein (AA 1-119) (His tag)



### Overview

Quantity:	100 μg
Target:	beta-2 Microglobulin (B2M)
Protein Characteristics:	AA 1-119
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This beta-2 Microglobulin protein is labelled with His tag.

## **Product Details**

Sequence:	Met1-Met119
Characteristics:	A DNA sequence encoding the Rat BMG/ $\beta$ 2-MG protein (P07151) (Met1-Met119) was expressed with a C-His.
Purity:	> 95 % as determined by reducing SDS-PAGE.

# **Target Details**

Target:	beta-2 Microglobulin (B2M)
Abstract:	B2M Products
Background:	Abbreviation: BMG,β2-MG
	Target Synonym: B2MG,Beta-2-Microglobulin,Beta 2 microglobin,Beta 2 microglobulin
	precursor,Beta chain of mhc class 1 proteins,Beta chain of MHC class I
	molecules,CDABP0092,Hdcma22p,IMD43

Background: β-2-Microglobulin (B2M) is a secreted protein with 1 Ig-like C1-type (immunoglobulin-like) domain which belongs to the beta-2-microglobulin family. B2M component of major histocompatibility complex (MHC) class I molecules, involved in the presentation of peptide antigens to the immune system. Polymers of beta 2-microglobulin can be found in tissues from patients on long-term hemodialysis. B2M is a protein found on the surface of many cells and plentiful on the surface of white blood cells. Serum B2M concentration is increased in renal diseases, various malignant diseases and some inflammatory and autoimmune disorders. B2M may adopt the fibrillar configuration of amyloid in certain pathologic states. The capacity to assemble into amyloid fibrils is concentration dependent. B2M has been shown as a marker for monitoring inflammatory disease activity and it appears likely to have a destructive role in amyloidosis-related arthritis. B2M might be involved in the OA (osteoarthritis) pathogenesis. Defects in B2M are the cause of hypercatabolic hypoproteinemia. Affected individuals show marked reduction in serum concentrations of immunoglobulin and albumin, probably due to rapid degradation. B2M could be a potential therapeutic target in ovarian cancer.

Molecular Weight:

Calculated MW: 12.98 kDa

Observed MW: 12 kDa

UniProt:

P07151

Pathways:

TCR Signaling, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune

**Effector Process** 

### **Application Details**

Restrictions:

For Research Use only

### Handling

Format:	Lyophilized
Buffer:	Lyophilized from sterile PBS, pH 7.4.  Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization.
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.  Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

		1.	
$\vdash$	land	lın	
	ıaııu	1111	L

Expiry Date:

12 months