# antibodies -online.com









Go to Product page

| $\sim$ |    |    |    |             |   |
|--------|----|----|----|-------------|---|
| ()     | VE | ۲۱ | /1 | $\triangle$ | Λ |


| Quantity:    | 100 tests                                 |  |
|--------------|---|--|
| Target:      | SIRPA                                     |  |
| Reactivity:  | Human                                     |  |
| Host:        | Mouse                                     |  |
| Clonality:   | Monoclonal                                |  |
| Conjugate:   | This SIRPA antibody is conjugated to FITC |  |
| Application: | Flow Cytometry (FACS)                     |  |


# **Product Details**

| Purpose:      | Anti-Hu CD172a FITC   |  |
|---------------|---|--|
| Immunogen:    | Kg-1a cell line   |  |
| Clone:        | 15-414  |  |
| Isotype:      | lgG2a   |  |
| Specificity:  | The mouse monoclonal antibody 15-414 recognizes en extracellular epitope of CD172a (SIRP alpha), an approximately 90 kDa transmembrane glycoprotein expressed on cells of myeloid origin and neurons. |  |
| Purification: | Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.         |  |

# **Target Details**

| Target:             | SIRPA  |  |
|---------------------|--|--|
| Alternative Name:   | CD172a (SIRPA Products)  |  |
| Background:         | Signal regulatory protein alpha,CD172a, the signal-regulatory protein alpha (SIRP alpha), also               |  |
|                     | known as SH2 domain-containing phosphatase substrate-1 (SHPS1), is a 75-110 kDa                              |  |
|                     | transmembrane glycoprotein expressed mainly on granulocytes, monocytes, macrophages,                         |  |
|                     | dendritic cells and neurons. Its extracellular ligand is CD47. CD172a serves as a substrate of               |  |
|                     | activated receptor tyrosine kinases and upon phosphorylation it recruits SH2 domain-                         |  |
|                     | containing tyrosine phosphatases, thereby regulating signal transduction processes related to                |  |
|                     | cell activation, transmigration and phagocytosis. CD172a is a specific marker of                             |  |
|                     | cardiomyocytes derived from human pluripotent stem cells and serves as a negative regulator                  |  |
|                     | of signaling and growth in myeloid progenitor cells.,PTPNS1, BIT, MFR, SIRPA, SHPS1                          |  |
| Gene ID:            | 140885   |  |
| UniProt:            | P78324   |  |
| Application Details |  |  |
| Application Notes:  | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µL reagent /               |  |
|                     | 100 $\mu L$ of whole blood or $10^6$ cells in a suspension. The content of a vial (0.4 ml) is sufficient for |  |
|                     | 100 tests.   |  |
| Restrictions:       | For Research Use only  |  |
| Handling            |  |  |
| Buffer:             | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide                                      |  |
| Preservative:       | Sodium azide   |  |
| Precaution of Use:  | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which                                |  |
|                     | should be handled by trained staff only.   |  |
| Storage:            | 4 °C   |  |
| Storage Comment:    | Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.                                     |  |





## **Flow Cytometry**

**Image 1.** Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD172a (15-414) FITC antibody (4  $\mu$ L reagent / 100  $\mu$ L of peripheral whole blood).

### **Flow Cytometry**

**Image 2.** Separation of human monocytes (red-filled) from lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD172a (15-414) FITC antibody (4  $\mu$ L reagent / 100  $\mu$ L of peripheral whole blood).