

Datasheet for ABIN7505957

**anti-CD79a antibody (AA 208-222) (FITC)**[Go to Product page](#)**3** Images

## Overview

Quantity:	100 tests
Target:	CD79a (CD79A)
Binding Specificity:	AA 208-222
Reactivity:	Human, Mouse, Rat, Cow, Pig, Chicken, Guinea Pig, Horse, Rabbit, Dog, Non-Human Primate
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD79a antibody is conjugated to FITC
Application:	Flow Cytometry (FACS)

## Product Details

Purpose:	Anti-Hu CD79a FITC
Immunogen:	Synthetic peptide corresponding to C terminal amino acids 208-222 of human CD79a
Clone:	HM47
Isotype:	IgG1 kappa
Specificity:	The mouse monoclonal antibody HM47 reacts with intracellular domain of CD79a (Ig alpha), a 40-45 kDa subunit of B cell antigen-specific receptor (BCR) and its early developmental forms.
Purification:	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

## Target Details


Target:	CD79a (CD79A)
Alternative Name:	CD79a ( <a href="#">CD79A Products</a> )
Background:	CD79a molecule,CD79a (Ig alpha, MB1) forms disulfide-linked heterodimer with CD79b (Ig beta). They both are transmembrane proteins with extended cytoplasmic domains containing immunoreceptor tyrosine activation motives (ITAMs), and together with cell surface immunoglobulin they constitute B-cell antigen-specific receptor (BCR). CD79a and b are the first components of BCR that are expressed developmentally. They appear on pro-B cells in association with the endoplasmic reticulum chaperone calnexin. Subsequently, in pre-B cells, CD79 heterodimer is associated with lambda5-VpreB surrogate immunoglobulin and later with antigen-specific surface immunoglobulins. At the plasma cell stage, CD79a is present as an intracellular component. CD79a/b complex interacts with Src-family tyrosine kinase Lyn, which phosphorylates its cytoplasmic ITAM motives to form docking sites for downstream signaling.,BCR alpha, Ig-alpha, MB-1, IGA
Gene ID:	973
UniProt:	<a href="#">P11912</a>
Pathways:	<a href="#">BCR Signaling</a>

## Application Details

Application Notes:	Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µL reagent / 100 µL of whole blood or 10 <sup>6</sup> cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. Intracellular staining.
Restrictions:	For Research Use only

## Handling

Buffer:	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.



### Flow Cytometry

**Image 1.** Flow cytometry intracellular staining pattern of human peripheral whole blood stained using anti-human CD79a (HM47) FITC antibody (4  $\mu$ L reagent / 100  $\mu$ L of peripheral whole blood).

### Flow Cytometry

**Image 2.** Flow cytometry multicolor intracellular staining pattern of human lymphocytes using anti-human CD79a (HM47) FITC antibody (4  $\mu$ L reagent / 100  $\mu$ L of peripheral whole blood) and anti-human CD19 (LT19) APC antibody (10  $\mu$ L reagent / 100  $\mu$ L of peripheral whole blood).

### Flow Cytometry

**Image 3.** Separation of human CD19 positive CD79a positive lymphocytes (red-filled) from neutrophil granulocytes (black-dashed) in flow cytometry analysis (intracellular staining) of human peripheral whole blood stained using anti-human CD79a (HM47) FITC antibody (4  $\mu$  L reagent / 100  $\mu$ L of peripheral whole blood).