antibodies -online.com









Overview

Quantity:	100 tests
Target:	SLC44A1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This SLC44A1 antibody is conjugated to PE
Application:	Flow Cytometry (FACS)

Product Details


Purpose:	Anti-Hu CD92 PE	
Immunogen:	MV4-11 acute myeloid leukemia cells	
Clone:	VIM15	
Isotype:	lgG1	
Specificity:	The mouse monoclonal antibody VIM15 recognizes an extracellular epitope of CD92, a 70 kDa transmembrane glycoprotein expressed mainly on human peripheral blood monocytes and neutrophils, and several myeloid and T-cell lines.	
Purification:	Purified antibody is conjugated with R-phycoerythrin (PE) under optimum conditions. Unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.	


Target Details

	01.04444	
Target:	SLC44A1	

Target Details

CD92 (SLC44A1 Products)	
Solute carrier family 44 member 1,CD92 is a 70 kDa protein with ten transmembrane domains,	
intracellular N and C teminus, and two glycosylated larger extracellular loops. In the C-terminal	
domain, there is an ITIM-like sequence. This protein seems to be a choline transporter	
responsible for delivery of choline into the immune cells, to make it accessible for phospholipid	
synthesis, as well as a regulator of immune cell signaling. It is expressed mainly on human	
peripheral blood monocytes and neutrophils, and several myeloid and T-cell lines. It can also be	
found on mast cells (but not eosinophils), and weakly on peripheral blood lymphocytes,	
fibroblasts, epithelial cells, and endothelial cells., CTL1, CHTL1, SLC44A1	
23446	
Q8WWI5	
Flow cytometry: The reagent is designed for analysis of human blood cells using 10 µL reagent	
/ 100 μL of whole blood or 10 6 cells in a suspension. The content of a vial (1 ml) is sufficient fo	
100 tests.	
For Research Use only	
Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide	
Sodium azide	
This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
should be handled by trained staff only.	
4 °C	
Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.	





Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral whole blood stained using anti-human CD92 (VIM15) PE antibody (10 μ L reagent / 100 μ L of peripheral whole blood).

Flow Cytometry

Image 2. Separation of human CD92 positive monocytes (red-filled) from lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD92 (VIM15) PE antibody (10 µL reagent / 100 µL of peripheral whole blood).