



[Go to Product page](#)

Datasheet for ABIN750870 anti-CUBN antibody (AA 51-150) (Biotin)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | CUBN |
| Binding Specificity: | AA 51-150 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CUBN antibody is conjugated to Biotin |
| Application: | ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human Cubilin |
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Predicted Reactivity: | Mouse,Rat |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|---|
| Target: | CUBN |
| Alternative Name: | Cubilin (CUBN Products) |

Target Details

| | |
|-------------|--|
| Background: | <p>Synonyms: IFCR, MGA1, gp280, Cubilin, 460 kDa receptor, Intestinal intrinsic factor receptor, Intrinsic factor-cobalamin receptor, Intrinsic factor-vitamin B12 receptor, CUBN</p> <p>Background: Cotransporter which plays a role in lipoprotein, vitamin and iron metabolism, by facilitating their uptake. Binds to ALB, MB, Kappa and lambda-light chains, TF, hemoglobin, GC, SCGB1A1, APOA1, high density lipoprotein, and the GIF-cobalamin complex. The binding of all ligands requires calcium. Serves as important transporter in several absorptive epithelia, including intestine, renal proximal tubules and embryonic yolk sac. Interaction with LRP2 mediates its trafficking throughout vesicles and facilitates the uptake of specific ligands like GC, hemoglobin, ALB, TF and SCGB1A1. Interaction with AMN controls its trafficking to the plasma membrane and facilitates endocytosis of ligands. May play an important role in the development of the peri-implantation embryo through internalization of APOA1 and cholesterol. Binds to LGALS3 at the maternal-fetal interface.</p> |
| Gene ID: | 8029 |
| UniProt: | O60494 |
| Pathways: | Metabolism of Steroid Hormones and Vitamin D , Lipid Metabolism |

Application Details

| | |
|--------------------|------------------------------------|
| Application Notes: | IHC-P 1:200-400 IHC-F 1:100-500 |
| Restrictions: | For Research Use only |

Handling

| | |
|--------------------|--|
| Format: | Liquid |
| Concentration: | 1 µg/µL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C for 12 months. |

Handling

Expiry Date: 12 months