antibodies - online.com







anti-ADCY4 antibody (AA 851-1077)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	ADCY4
Binding Specificity:	AA 851-1077
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ADCY4 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ADCY4
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	ADCY4	

Target Details

rarget Details	
Alternative Name:	ADCY4 (ADCY4 Products)
Background:	Synonyms: AC4, Adenylate cyclase 4, adenylate cyclase type 4, Adenylate cyclase type IV,
	Adenylyl cyclase, ATP pyrophosphate lyase 4, ATP pyrophosphate lyase, ADCY4_HUMAN.
	Background: The ADCY4 gene encodes a member of the family of adenylate cyclases, which
	are membrane-associated enzymes that catalyze the formation of the secondary messenger
	cyclic adenosine monophosphate (cAMP). Mouse studies show that adenylate cyclase 4, along
	with adenylate cyclases 2 and 3, is expressed in olfactory cilia, suggesting that several differen
	adenylate cyclases may couple to olfactory receptors and that there may be multiple receptor-
	mediated mechanisms for the generation of cAMP signals. Alternative splicing results in
	transcript variants.
Gene ID:	196883
Pathways:	EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Thyroid Hormone Synthesis, cAMP
	Metabolic Process, Myometrial Relaxation and Contraction, G-protein mediated Events,
	Interaction of EGFR with phospholipase C-gamma
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.

Handling

Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months