# antibodies -online.com







# anti-ADCY4 antibody (AA 851-1077) (Biotin)



( )	1/0	r\ / I	014	
( )	ve	I V I	-v	V

Quantity:	100 μL	
Target:	ADCY4	
Binding Specificity:	AA 851-1077	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ADCY4 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human ADCY4
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

### **Target Details**

Target:	ADCY4
Alternative Name:	ADCY4 (ADCY4 Products)

## **Target Details**

rarget Details	
Background:	Synonyms: AC4, Adenylate cyclase 4, adenylate cyclase type 4, Adenylate cyclase type IV, Adenylyl cyclase, ATP pyrophosphate lyase 4, ATP pyrophosphate lyase, ADCY4_HUMAN. Background: The ADCY4 gene encodes a member of the family of adenylate cyclases, which are membrane-associated enzymes that catalyze the formation of the secondary messenger cyclic adenosine monophosphate (cAMP). Mouse studies show that adenylate cyclase 4, along with adenylate cyclases 2 and 3, is expressed in olfactory cilia, suggesting that several different adenylate cyclases may couple to olfactory receptors and that there may be multiple receptor-mediated mechanisms for the generation of cAMP signals. Alternative splicing results in transcript variants.
Gene ID:	196883
Pathways:	EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Thyroid Hormone Synthesis, cAMP Metabolic Process, Myometrial Relaxation and Contraction, G-protein mediated Events, Interaction of EGFR with phospholipase C-gamma
Application Details	
Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

handled by trained staff only.

Store at -20°C for 12 months.

-20 °C

12 months

Storage:

Expiry Date:

Storage Comment: