antibodies .- online.com





anti-SLC27A2 antibody (AA 401-500) (Cy5.5)



()	11/0	K\ /	iew
	\cup	'I V/I	$I \cap VV$

Quantity:	100 μL
Target:	SLC27A2
Binding Specificity:	AA 401-500
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLC27A2 antibody is conjugated to Cy5.5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SLC27A2/ACSVL1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	SLC27A2
Alternative Name:	SLC27A2/ACSVL1 (SLC27A2 Products)

Target Details

Background:	Synonyms: VLCS, FATP2, VLACS, ACSVL1, FACVL1, hFACVL1, HsT17226, Very long-chain acyl-
	CoA synthetase, Fatty acid transport protein 2, FATP-2, Fatty-acid-coenzyme A ligase, very long
	chain 1, Long-chain-fatty-acidCoA ligase, Solute carrier family 27 member 2, THCA-CoA ligase,
	Very long-chain-fatty-acid-CoA ligase, SLC27A2
	Background: Acyl-CoA synthetase probably involved in bile acid metabolism. Proposed to
	activate C27 precurors of bile acids to their CoA thioesters derivatives before side chain
	cleavage via peroxisomal beta-oxidation occurs. In vitro, activates 3-alpha,7-alpha,12-alpha-
	trihydroxy-5-beta-cholestanate (THCA), the C27 precursor of cholic acid deriving from the de
	novo synthesis from cholesterol. Does not utilize C24 bile acids as substrates. In vitro, also
	activates long- and branched-chain fatty acids and may have additional roles in fatty acid
	metabolism. May be involved in translocation of long-chain fatty acids (LFCA) across
	membranes (By similarity).
Gene ID:	11001
UniProt:	014975
Pathways:	Monocarboxylic Acid Catabolic Process, SARS-CoV-2 Protein Interactome
Application Dataila	
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months