Datasheet for ABIN751402 anti-SLC27A2 antibody (AA 401-500) (HRP)

-online.com antibodies



Overview

Quantity:	100 µL
Target:	SLC27A2
Binding Specificity:	AA 401-500
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLC27A2 antibody is conjugated to HRP
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SLC27A2/ACSVL1
Isotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.
Target Details	
Target:	SLC27A2
Alternative Name:	SLC27A2/ACSVL1 (SLC27A2 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN751402 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

yms: VLCS, FATP2, VLACS, ACSVL1, FACVL1, hFACVL1, HsT17226, Very long-chain acyl- ynthetase, Fatty acid transport protein 2, FATP-2, Fatty-acid-coenzyme A ligase, very long- 1, Long-chain-fatty-acidCoA ligase, Solute carrier family 27 member 2, THCA-CoA ligase, ong-chain-fatty-acid-CoA ligase, SLC27A2 round: Acyl-CoA synthetase probably involved in bile acid metabolism. Proposed to ce C27 precurors of bile acids to their CoA thioesters derivatives before side chain ge via peroxisomal beta-oxidation occurs. In vitro, activates 3-alpha,7-alpha,12-alpha- oxy-5-beta-cholestanate (THCA), the C27 precursor of cholic acid deriving from the de
I, Long-chain-fatty-acidCoA ligase, Solute carrier family 27 member 2, THCA-CoA ligase, ong-chain-fatty-acid-CoA ligase, SLC27A2 round: Acyl-CoA synthetase probably involved in bile acid metabolism. Proposed to ce C27 precurors of bile acids to their CoA thioesters derivatives before side chain ge via peroxisomal beta-oxidation occurs. In vitro, activates 3-alpha,7-alpha,12-alpha-
ong-chain-fatty-acid-CoA ligase, SLC27A2 round: Acyl-CoA synthetase probably involved in bile acid metabolism. Proposed to ce C27 precurors of bile acids to their CoA thioesters derivatives before side chain ge via peroxisomal beta-oxidation occurs. In vitro, activates 3-alpha,7-alpha,12-alpha-
round: Acyl-CoA synthetase probably involved in bile acid metabolism. Proposed to the C27 precurors of bile acids to their CoA thioesters derivatives before side chain ge via peroxisomal beta-oxidation occurs. In vitro, activates 3-alpha,7-alpha,12-alpha-
e C27 precurors of bile acids to their CoA thioesters derivatives before side chain ge via peroxisomal beta-oxidation occurs. In vitro, activates 3-alpha,7-alpha,12-alpha-
ge via peroxisomal beta-oxidation occurs. In vitro, activates 3-alpha,7-alpha,12-alpha-
oxy-5-beta-cholestanate (THCA), the C27 precursor of cholic acid deriving from the de
ynthesis from cholesterol. Does not utilize C24 bile acids as substrates. In vitro, also
es long- and branched-chain fatty acids and may have additional roles in fatty acid
olism. May be involved in translocation of long-chain fatty acids (LFCA) across
ranes (By similarity).
5
carboxylic Acid Catabolic Process, SARS-CoV-2 Protein Interactome
300-5000
1:200-400
1:100-500
search Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN751402 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Handling	
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months