

Datasheet for ABIN751560 anti-IDH2 antibody (AA 251-350) (Biotin)



Go to Product page

| Overview | |
|-----------------------|---|
| Quantity: | 100 μL |
| Target: | IDH2 |
| Binding Specificity: | AA 251-350 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This IDH2 antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |
| Product Details | |
| Immunogen: | KLH conjugated synthetic peptide derived from human IDH2 |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Predicted Reactivity: | Dog,Horse |
| Purification: | Purified by Protein A. |
| Target Details | |
| Target: | IDH2 |
| Alternative Name: | IDH2 (IDH2 Products) |

Target Details

Expiry Date:

12 months

| rarget Details | |
|---------------------|--|
| Background: | Synonyms: IDH, IDP, IDHM, IDPM, ICD-M, D2HGA2, mNADP-IDH, Isocitrate dehydrogenase [NADP], mitochondrial, NADP(+)-specific ICDH, Oxalosuccinate decarboxylase, IDH2 Background: Plays a role in intermediary metabolism and energy production. It may tightly associate or interact with the pyruvate dehydrogenase complex. |
| Gene ID: | 3418 |
| UniProt: | P48735 |
| Pathways: | Warburg Effect |
| Application Details | |
| Application Notes: | WB 1:300-5000 |
| | IHC-P 1:200-400 |
| | IHC-F 1:100-500 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C for 12 months. |
| Evening Deter | 10 months |