## antibodies .- online.com





## anti-MT-ND5 antibody (AA 451-550) (Cy3)



Go to Product page

( )	11/0	K\ /	iew
	$\cup$	'I V/I	$I \cap VV$

Quantity:	100 μL
Target:	MT-ND5
Binding Specificity:	AA 451-550
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MT-ND5 antibody is conjugated to Cy3
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human MT-ND5
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Purification:	Purified by Protein A.

## **Target Details**

Target:	MT-ND5
Alternative Name:	MT-ND5 (MT-ND5 Products)
Background:	Synonyms: MTND5, NADH-ubiquinone oxidoreductase chain 5, NADH dehydrogenase subunit

Expiry Date:

12 months

Target Details	
	5, MT-ND5, NADH5, ND5
	Background: Core subunit of the mitochondrial membrane respiratory chain NADH
	dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for
	catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain.
	The immediate electron acceptor for the enzyme is believed to be ubiquinone (By similarity).
Gene ID:	4540
UniProt:	P03915
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.