antibodies.com

Datasheet for ABIN751755 anti-NDUFV2 antibody (AA 101-200) (Biotin)



\sim			
Οv	Dr\	110	1/ / /
Οv	CIV	VIC	- V V

Quantity:	100 µL	
Target:	NDUFV2	
Binding Specificity:	AA 101-200	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This NDUFV2 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NDUFV2
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NDUFV2
Alternative Name:	NDUFV2 (NDUFV2 Products)
Background:	Synonyms: 24 kDa subunit of Complex I, CI-24k, NADH dehydrogenase ubiquinone flavoprotein

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN751755 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

	2, 24 kDa, NADH ubiquinone oxidoreductase 24 kDa subunit, Ubiquinoneflavoprotein 2,
	mitochondrial precursor.
	Background: The NADH-ubiquinone oxidoreductase complex (complex I) of the mitochondrial
	respiratory chain catalyzes the transfer of electrons from NADH to ubiquinone, and consists of
	at least 43 subunits. The complex is located in the inner mitochondrial membrane. This gene
	encodes the 24 kDa subunit of complex I, and is involved in electron transfer. Mutations in this
	gene are implicated in Parkinson's disease, bipolar disorder, schizophrenia, and have been
	found in one case of early onset hypertrophic cardiomyopathy and encephalopathy. A non-
	transcribed pseudogene of this locus is found on chromosome 19. [provided by RefSeq, Oct
	2009].
Gene ID:	4729
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months