antibodies -online.com







anti-PDP antibody (AA 51-150)



Overview

Quantity:	100 μL
Target:	PDP
Binding Specificity:	AA 51-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PDP antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PPM2C
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Horse,Rabbit
Purification:	Purified by Protein A.
Target Details	

rarget Details

Target:	PDP
Alternative Name:	PPM2C (PDP Products)

Target Details

Expiry Date:

12 months

9	
Background:	Synonyms: PDH, PDP 1, PDP, PDP1, PDPC 1, PDPC, Protein phosphatase 2C, protein
	phosphatase 2C, magnesium dependent, catalytic subunit, Pyruvate dehydrogenase
	phosphatase, catalytic subunit 1, PDP1_HUMAN.
	Background: PPM2C is a protein similar to members of the protein phosphatase 2C (PP2C)
	family. It is a catalytic subunit of the mitochondrial matrix phosphatase and is involved in
	reactivation of the pyruvate dehydrogenase complex.
Gene ID:	54704
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.