

Datasheet for ABIN7519857

CXCL1 Protein



Go to Product page

_			
()	V/C	rv	٨/

Quantity:	10 μg
Target:	CXCL1
Origin:	Mouse
Source:	Yeast (Pichia pastoris)
Protein Type:	Recombinant

Product Details

Purpose:	Recombinant Mouse CXCL1/GRO-alpha Protein
Sequence:	RLATGAPIAN ELRCQCLQTM AGIHLKNIQS LKVLPSGPHC TQTEVIATLK NGREACLDPE APLVQKIVQK MLKGVPK
Specificity:	Arg20-Lys96
Purity:	> 95 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	<0.1EU/µg

Target Details

Target:	CXCL1	
Alternative Name:	CXCL1/GRO-alpha (CXCL1 Products)	
Background:	Description: This antimicrobial protein belongs a member of the CXC subfamily of chemokines.	
	The encoded protein is a secreted growth factor that signals through the G-protein coupled	
	receptor, CXC receptor 2. This protein plays a role in inflammation and as a chemoattractant for	

Target Details

neutrophils. Aberrant expression of this protein is associated with the growth and progression		
of certain tumors. A naturally occurring processed form of this protein has increased		
chemotactic activity. Alternate splicing results in coding and non-coding variants of this gene. A		
pseudogene of this gene is found on chromosome 4.		
Name: FSP, GRO1, GROa, MGSA, MGSA-a, NAP-3, SCYB1,CXCL1		

Gene ID:

14825

UniProt:

P12850

Pathways:

Autophagy

Application Details

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile
	distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is
	recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 %
	Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.
Buffer:	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.
Storage:	-20 °C,-80 °C
Storage Comment:	Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein
	solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.