

Datasheet for ABIN7519915

CST1 Protein (His tag)



Overview

Quantity:	500 μg
Target:	CST1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This CST1 protein is labelled with His tag.

Product Details

Purpose:	Active Recombinant Human Cystatin-SN/CST1 Protein
Sequence:	WSPKEEDRII PGGIYNADLN DEWVQRALHF AISEYNKATK DDYYRRPLRV LRARQQTVGG VNYFFDVEVG RTICTKSQPN LDTCAFHEQP ELQKKQLCSF EIYEVPWENR RSLVKSRCQE S
Specificity:	Trp21-Ser141
Purity:	> 97 % by SDS-PAGE.
Sterility:	0.22 μm filtered
Endotoxin Level:	< 0.1 EU/µg of the protein by LAL method.
Biological Activity Comment:	Measured by its ability to inhibit active Cathepsin L cleavage of a fluorogenic peptide substrate Z-LR-AMC. The IC50 value is <11.0 nM.

Target Details

rarget Details	
Target:	CST1
Alternative Name:	Cystatin-SN/CST1 (CST1 Products)
Background:	Description: The cystatin superfamily encompasses proteins that contain multiple cystatin-like
	sequences. Some of the members are active cysteine protease inhibitors, while others have los
	or perhaps never acquired this inhibitory activity. There are three inhibitory families in the
	superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The
	type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human
	fluids and secretions, where they appear to provide protective functions. The cystatin locus on
	chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This
	gene is located in the cystatin locus and encodes a cysteine proteinase inhibitor found in saliva
	tears, urine, and seminal fluid.
	Name: CST1
Gene ID:	1469
UniProt:	P01037
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile
	distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is
	recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 %
	Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.
Buffer:	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.
Storage:	-20 °C,-80 °C
Storage Comment:	Store the lyophilized protein at -20°C to -80 °C for long term.
	After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1
	week.