

## Datasheet for ABIN7519930

# **MYDGF Protein (His tag)**



#### Overview

Quantity:	50 μg
Target:	MYDGF (D17Wsu104e)
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This MYDGF protein is labelled with His tag.

### **Product Details**

Purpose:	Recombinant Human MYDGF Protein
Sequence:	VSEPTTVAFD VRPGGVVHSF SHNVGPGDKY TCMFTYASQG GTNEQWQMSL GTSEDHQHFT CTIWRPQGKS YLYFTQFKAE VRGAEIEYAM AYSKAAFERE SDVPLKTEEF EVTKTAVAHR
	PGAFKAELSK LVIVAKASRT EL
Specificity:	Val32-Leu173
Purity:	> 97 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	< 0.1EU/µg

## Target Details

Target:	MYDGF (D17Wsu104e)
Alternative Name:	MYDGF (D17Wsu104e Products)

Target Details	
Background:	Description: Myeloid-Derived Growth Factor, or MYDGF, is a Bone marrow-derived monocyte
	protein, and it is correlated with enhanced metabolic activity, suppression of apoptosis, and
	stimulation of cell proliferation . MYDGF is expressed predominantly in inflammatory cells, such
	as monocytes and macrophages . Up-regulation of MYDGF expression was also found during
	adipocyte differentiation . Expression of MYDGF was induced in the circulation and heart tissue
	after myocardial infarction. It promotes cardiac myocyte survival by stimulating endothelial cell
	proliferation through a MAPK1/3-, STAT3- and CCND1-mediated signaling pathway, and inhibits
	cardiac myocyte apoptosis in a PI3K/AKT-dependent signaling pathway . MYDGF was found
	over-expressed in approximately two-thirds of Hepatocellular Carcinoma (HCC) tissues, and its
	expression was significantly positively correlated with that of alpha-fetoprotein (AFP) .
	Name: IL25, IL27, SF20, IL27w, C19orf10, R33729_1, EUROIMAGE1875335,MYDGF
Gene ID:	56005
UniProt:	Q969H8
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile
	distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is

Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.
Concentration:	1.46 mg/mL
Buffer:	Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.
Storage:	-20 °C,-80 °C
Storage Comment:	Store the lyophilized protein at -20°C to -80°C for 12 months. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.