

[Go to Product page](#)

## Datasheet for ABIN7520009 FGFR2 Protein (Fc Tag,His tag)

### Overview

|                               |   |
|-------------------------------|---|
| Quantity:                     | 50 µg   |
| Target:                       | FGFR2   |
| Origin:                       | Human   |
| Source:                       | HEK-293 Cells                                       |
| Protein Type:                 | Recombinant   |
| Biological Activity:          | Active  |
| Purification tag / Conjugate: | This FGFR2 protein is labelled with Fc Tag,His tag. |

### Product Details

|                              |   |
|------------------------------|---|
| Purpose:                     | Active Recombinant Human FGFR-2/KGFR/CD332 Protein  |
| Sequence:                    | RPSFSLVEDT TLEPEEPPTK YQISQPEVYV AAPGESLEVR CLLKDAAVIS WTKDGVHLGP<br>NNRTVLIGEY LQIKGATPRD SGLYACTASR TVDSETWYFM VNVTDAISSG DDEDDTDGAE<br>DFVSENSNNK RPYWTNTEK MEKRLHAVPA ANTVKFRCPA GGNPMPTMRW LKNGKEFKQE<br>HRIGGYKVRN QHWSLIMESV VPSDKGNYTC VVENEYGSIN HTYHLDVVER SPHRPILQAG<br>LPANASTVVG GDVEFVCKVY SDAQPHIQWI KHVEKNGSKY GPDGLPYLKV LKAAGVNTTD<br>KEIEVLYIRN VTFEDAGEYT CLAGNSIGIS FHSAWLTVLP APGREKEITA SPDYLE |
| Specificity:                 | Arg22-Glu377  |
| Purity:                      | > 97 % by SDS-PAGE.   |
| Sterility:                   | 0.22 µm filtered  |
| Endotoxin Level:             | < 0.1 EU/µg of the protein by LAL method.   |
| Biological Activity Comment: | 1.Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human FGF1   |

## Product Details

at 5 µg/mL (100 µL/well) can bind Recombinant Human FGFR2 with a linear range of 0.8-2.5 µg/mL.<sup>[2]</sup> Measured by its ability to inhibit FGF-acidic dependent proliferation of Balb/c 3T3 mouse fibroblasts. The ED<sub>50</sub> for this effect is typically 0.256-0.991 ng/mL.

## Target Details

|                   |  |
|-------------------|--|
| Target:           | FGFR2  |
| Alternative Name: | FGFR-2/KGFR/CD332 ( <a href="#">FGFR2 Products</a> )   |
| Background:       | <p>Description: The protein is a member of the fibroblast growth factor receptor family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member is a high-affinity receptor for acidic, basic and/or keratinocyte growth factor, depending on the isoform. Mutations in this gene are associated with Crouzon syndrome, Pfeiffer syndrome, Craniosynostosis, Apert syndrome, Jackson-Weiss syndrome, Beare-Stevenson cutis gyrata syndrome, Saethre-Chotzen syndrome, and syndromic craniosynostosis.</p> <p>Name: BBDS,BEK,BFR-1,CD332,CEK3,CFD1,ECT1,JWS,K-SAM,KGFR,TK14,TK25,FGFR2</p> |
| Gene ID:          | 2263   |
| UniProt:          | <a href="#">P21802-1</a>   |
| Pathways:         | <a href="#">RTK Signaling</a> , <a href="#">Fc-epsilon Receptor Signaling Pathway</a> , <a href="#">EGFR Signaling Pathway</a> , <a href="#">Neurotrophin Signaling Pathway</a> , <a href="#">Regulation of Muscle Cell Differentiation</a> , <a href="#">Skeletal Muscle Fiber Development</a> , <a href="#">Growth Factor Binding</a>  |

## Application Details

|                 |   |
|-----------------|---|
| Restrictions:   | For Research Use only   |
| Handling        |   |
| Format:         | Lyophilized   |
| Reconstitution: | Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile |

## Handling

---

distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

---

|         |  |
|---------|--|
| Buffer: | Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. |
|---------|--|

---

|          |                |
|----------|----------------|
| Storage: | -20 °C, -80 °C |
|----------|----------------|

---

|                  |  |
|------------------|--|
| Storage Comment: | <p>Store the lyophilized protein at -20°C to -80 °C for long term.</p> <p>After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.</p> |
|------------------|--|