antibodies .- online.com







FKBP1A Protein



Overview

Quantity:	10 μg
Target:	FKBP1A
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant

Product Details

Purpose:	Recombinant Human FKBP1A/FKBP12 Protein
Sequence:	MGVQVETISP GDGRTFPKRG QTCVVHYTGM LEDGKKFDSS RDRNKPFKFM LGKQEVIRGW EEGVAQMSVG QRAKLTISPD YAYGATGHPG IIPPHATLVF DVELLKLE
Specificity:	Met1-Glu108
Purity:	> 90 % by SDS-PAGE.
Sterility:	0.22 µm filtered
Endotoxin Level:	< 0.1 EU/µg of the protein by LAL method.

Target Details

Target:	FKBP1A
Alternative Name:	FKBP1A/FKBP12 (FKBP1A Products)
Background:	Description: The protein is a member of the immunophilin protein family, which play a role in immunoregulation and basic cellular processes involving protein folding and trafficking. The
	protein is a cis-trans prolyl isomerase that binds the immunosuppressants FK506 and

rapamycin. It interacts	with several intracellular signal transduction proteins including type I
TGF-beta receptor. It a	lso interacts with multiple intracellular calcium release channels, and
coordinates multi-prot	ein complex formation of the tetrameric skeletal muscle ryanodine
receptor.	

Name: FKBP-12, FKBP-1A, FKBP1, FKBP12, PKC12, PKC12, PPIASE, FKBP1A, FKBP-1A, FKBP1, FKBP12, PKC12, PPIASE

Gene ID:	2280

UniProt: P62942

Pathways: Negative Regulation of Transporter Activity, Methionine Biosynthetic Process

Application Details

Restrictions: For Research Use only

Handling

Format:	Lyophilized	
Reconstitution:	Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1 % BSA, 5 % HSA, 10 % FBS or 5 % Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.	
Buffer:	Lyophilized from a 0.22 µm filtered solution of 20 mM Tris, 150 mM NaCl, pH 8.0.	
Storage:	-20 °C,-80 °C	
Storage Comment:	Store the lyophilized protein at -20°C to -80 °C for long term. After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.	